

OPEN QUANTUM SYSTEMS

A. Isar*, A. Sandulescu*, H. Scutaru*, E. Stefanescu* and W. Scheid#

**Department of Theoretical Physics, Institute of Atomic Physics,
Bucharest-Magurele, Romania*

*#Institut für Theoretische Physik der Justus-Liebig- Universität,
Giessen, Germany*

ABSTRACT

The damping of the harmonic oscillator is studied in the framework of the Lindblad theory for open quantum systems. A generalization of the fundamental constraints on quantum mechanical diffusion coefficients which appear in the master equation for the damped quantum oscillator is presented; the Schrödinger, Heisenberg and Weyl-Wigner-Moyal representations of the Lindblad equation are given explicitly. On the basis of these representations it is shown that various master equations for the damped quantum oscillator used in the literature are particular cases of the Lindblad equation and that not all of these equations are satisfying the constraints on quantum mechanical diffusion coefficients. Analytical expressions for the first two moments of coordinate and momentum are obtained by using the characteristic function of the Lindblad master equation. The master equation is transformed into Fokker-Planck equations for quasiprobability distributions and a comparative study is made for the Glauber P representation, the antinormal ordering Q representation and the Wigner W representation. The density matrix is represented via a generating function, which is obtained by solving a time-dependent linear partial differential equation derived from the master equation. Illustrative examples for specific initial conditions of the density matrix are provided. The solution of the master equation in the Weyl-Wigner-Moyal representation is of Gaussian type if the initial form of the Wigner function is taken to be a Gaussian corresponding (for example) to a coherent wavefunction. The damped harmonic oscillator is applied for the description of the charge equilibration mode observed in deep inelastic reactions. For a system consisting of two harmonic oscillators the time dependence of expectation values, Wigner function and Weyl operator are obtained and discussed. In addition models for the damping of the angular momentum are studied. Using this theory to the quantum tunneling through the nuclear barrier, besides Gamow's transitions with energy conservation, additional transitions with energy loss, are found. The tunneling spectrum is obtained as a function of the barrier characteristics. When this theory is used to the resonant atom-field interaction, new optical equations describing the coupling through the environment of the atomic observables are obtained. With these equations, some characteristics of the laser radiation absorption spectrum and optical bistability are described.

1. Introduction

In the last two decades, more and more interest arose about the problem of dissipation in quantum mechanics, i.e. the consistent description of open quantum systems [1-7]. The quantum description of dissipation is important in many different areas of physics. In quantum optics we mention the quantum theory of lasers and photon detection. There are some directions in the theory of atomic nucleus in which dissipative processes play a basic role. In this sense we mention the nuclear fission, giant resonances and deep inelastic collisions of heavy ions. Dissipative processes often occur also in many body or field-theoretical systems.

The irreversible, dissipative behaviour of the vast majority of physical phenomena comes into an evident contradiction with the reversible nature of our basic models. The very restrictive principles of conservative and isolated systems are unable to deal with more complicated situations which are determined by the features of open systems.

The fundamental quantum dynamical laws are of the reversible type. The dynamics of a closed system is governed by the Hamiltonian that represents its total energy and which is a constant of motion. In this way the paradox of irreversibility arises: the reversibility of microscopic dynamics contrasting with the irreversibility of the macroscopic behaviour we are trying to deduce from it.

One way to solve this paradox of irreversibility is to use models to which Hamiltonian dynamics and Liouville's theorem do not apply, but the irreversible behaviour is clearly present even in the microscopic dynamical description. The reason for replacing Hamiltonian dynamics and Liouville's theorem is that no system is truly isolated, being subject to uncontrollable random influences from outside. For this reason these models are called open systems. There are two ways of treating quantitatively their interaction with the outside. One way is to introduce specific stochastic assumptions to simulate this interaction, the other one is to treat the open systems with the usual laws of dynamics, by regarding them as subsystems of larger systems which are closed (i.e. which obey the usual laws of dynamics with a well-defined Hamiltonian). The dissipation arises in general from the subsystem interactions with a larger system, often referred to as the reservoir or bath. The first of these two approaches has been used for the study of steady state transport processes in systems obeying classical mechanics. The second of the two approaches has

been mainly used in quantum mechanics. The main general result [2,8-10] is that under certain conditions the time evolution of an open system can be described by a dynamical semigroup $\Phi_t(t \geq 0)$. For a closed finite system with Liouville operator the evolution operator is not restricted to nonnegative t . The importance of the dynamical semigroup concept is that it generalizes the evolution operator to open systems, for which there is no proper Liouville operator and no Φ_t for negative t . The mathematical theory of dynamical semigroups has been developed in [2,11-15].

In Sect.2 the notion of the quantum dynamical semigroup is defined using the concept of a completely positive map [14]. The Lindblad formalism replaces the dynamical group uniquely determined by its generator, which is the Hamiltonian operator of the system, by the completely positive dynamical semigroup with bounded generators. Then the general form of Markovian quantum mechanical master equation is obtained.

The quantum mechanics of the one-dimensional damped harmonic oscillator represents a fundamental theoretical problem with applications in different domains of quantum optics, solid state physics, molecular and nuclear physics. In the present review paper the quantum harmonic oscillator is treated in the Lindblad axiomatic formalism of quantum dynamical semigroups. In Sect.3 we give the fundamental constraints on quantum mechanical diffusion coefficients which appear in the corresponding master equations [16,17]. On the basis of different representations it is shown that various master equations for the damped quantum oscillator used in the literature for the description of the damped collective modes in deep inelastic collisions or in quantum optics are particular cases of the Lindblad equation and that not all of these equations in literature are satisfying the constraints on quantum mechanical diffusion coefficients. In 3.1, by using the Heisenberg representation, explicit expressions of the mean values and variances are given [17]. In 3.2 we solve the master equation with the characteristic function [18]. This function is found as a solution of a corresponding partial differential equation. By this method one can derive explicit formulae for the centroids and variances and, in general, for moments of any order. In 3.3 we explore the applicability of quasiprobability distributions to the Lindblad theory [19]. The methods of quasiprobabilities have provided technical tools of great power for the statistical description of microscopic systems formulated in terms of the density operator. In the present review the master equation of the one-dimensional

damped harmonic oscillator is transformed into Fokker-Planck equations for the Glauber P , antinormal Q and Wigner W quasiprobability distributions associated with the density operator. A comparative study of these representations is made. The resulting equations are solved by standard methods and observables directly calculated as correlations of these distribution functions. We solve also the Fokker-Planck equations for the steady state and show that variances found from the P, Q and W distributions are the same [19]. In 3.4 we study the time evolution of the density matrix that follows from the master equation of the damped harmonic oscillator [20]. We calculate the physically relevant solutions (including both diagonal and off-diagonal matrix elements) of the master equation by applying the method of generating function. This means that we represent the density matrix with a generating function which is the solution of a time-dependent partial differential equation of second order, derived from the master equation of the damped harmonic oscillator. We discuss stationary solutions of the generating function and derive the Bose-Einstein density matrix as example. Then, formulas for the time evolution of the density matrix are presented and illustrative examples for specific initial conditions provided. The same method of generating function was already used by Jang [21] who studied the damping of a collective degree of freedom coupled to a bosonic reservoir at finite temperature with a second order RPA master equation in the collective subspace. In 3.5 the master equation is given in the Weyl-Wigner-Moyal representation [17]. We show that the solution of the Lindblad equation in this representation is of Gaussian type if the initial form of the Wigner function is taken to be a Gaussian corresponding to a coherent wavefunction.

In Sect.4 we give some applications to nuclear equilibration processes. In 4.1 the charge equilibration mode is treated by a damped harmonic oscillator in the framework of the Lindblad theory [17]. It is shown that the centroids and variances of the charge equilibration mode observed in deep inelastic reactions can be well described by the corresponding overdamped solutions [18,22]. In 4.2 we treat the damping of the proton and neutron asymmetry degrees of freedom with the method of Lindblad by studying the damping of two coupled oscillators [23]. We present the equation of motion of the open quantum system of two oscillators in the Heisenberg picture and we derive the time dependence of the expectation values of the coordinates and momenta and their variances. We discuss also the connection with the Wigner function and Weyl operator. We demonstrate the

time dependence of the various quantities for a simplified version of the model, where the decay constants can be calculated analytically.

In Sect.5 simple models for the damping of the angular momentum are studied in the framework of the Lindblad theory of open systems [24,25]. We assume that the system is opened by the generators of the proper Lorentz group and the Heisenberg group.

In Sect.6 we study the quantum tunneling [26]. Considering the tunneling operator between the localized states, we solve Lindblad master equation in the second order approximation of the theory of perturbations. We find that diffusion leads to additional tunneling processes around the Q -value, while friction determines only transitions at lower energies. Generally, this result is contrary to the majority of the previous studies [27,28] where, usually, it was concluded that the dissipation should decrease the tunneling rate.

In Sect.7 we study the resonant atom-field interaction [29]. From Lindblad's theorem, instead of the conventional optical Bloch equations, where only decay processes appear, we obtain a more general system of equations where the atomic observables are coupled through the environment. Based on the new equations, experimental characteristics of the electromagnetic field absorption spectrum and of the optical bistability are described. At the same time, we predict that in some conditions, an energy transfer from the dissipative environment to the coherent electromagnetic field appears.

The concluding summary is given in Sect.8.

2. Lindblad theory for open quantum systems

The standard quantum mechanics is Hamiltonian. The time evolution of a closed physical system is given by a dynamical group U_t which is uniquely determined by its generator H , which is the Hamiltonian operator of the system. The action of the dynamical group U_t on any density matrix ρ from the set $\mathcal{D}(\mathcal{H})$ of all density matrices of the quantum system, whose corresponding Hilbert space is $\mathcal{H}$, is defined by

$$\rho(t) = U_t(\rho) = e^{-\frac{i}{\hbar}Ht} \rho e^{\frac{i}{\hbar}Ht}$$

for all $t \in (-\infty, \infty)$. We remind that, according to von Neumann, density operators $\rho \in \mathcal{D}(\mathcal{H})$ are trace class ($\text{Tr}\rho < \infty$), self-adjoint ($\rho^+ = \rho$), positive ($\rho > 0$) operators with $\text{Tr}\rho = 1$. All these properties are conserved by the time evolution defined by U_t .

In the case of open quantum systems the main difficulty consists in finding such time evolutions Φ_t for density operators $\rho(t) = \Phi_t(\rho)$ which preserve the von Neumann

conditions for all times. From the last requirement it follows that Φ_t must have the following properties:

- (i) $\Phi_t(\lambda_1 \rho_1 + \lambda_2 \rho_2) = \lambda_1 \Phi_t(\rho_1) + \lambda_2 \Phi_t(\rho_2); \lambda_1, \lambda_2 \geq 0$ with $\lambda_1 + \lambda_2 = 1$,
- (ii) $\Phi_t(\rho^+) = \Phi_t(\rho)^+$,
- (iii) $\Phi_t(\rho) > 0$,
- (iv) $\text{Tr} \Phi_t(\rho) = 1$.

But these conditions are not restrictive enough in order to give a complete description of the mappings Φ_t as in the case of the time evolutions U_t for closed systems. Even in this last case one has to impose other restrictions to U_t , namely, it must be a group $U_{t+s} = U_t U_s$. Also, it is evident that in this case $U_0(\rho) = \rho$ and $U_t(\rho) \rightarrow \rho$ in the trace norm when $t \rightarrow 0$. For the dual group $\tilde{U}_t$ acting on the observables $A \in \mathcal{B}(\mathcal{H})$, i.e. on the bounded operators on $\mathcal{H}$, we have

$$\tilde{U}_t(A) = e^{\frac{i}{\hbar} H t} A e^{-\frac{i}{\hbar} H t}.$$

Then $\tilde{U}_t(AB) = \tilde{U}_t(A) \tilde{U}_t(B)$ and $\tilde{U}_t(I) = I$, where I denotes the identity operator on $\mathcal{H}$. Also, $\tilde{U}_t(A) \rightarrow A$ ultraweakly when $t \rightarrow 0$ and $\tilde{U}_t$ is an ultraweakly continuous mapping [2,9,13,14,30]. These mappings have a strong positivity property called complete positivity:

$$\sum_{i,j} B_i^+ \tilde{U}_t(A_i^+ A_j) B_j \geq 0, \quad A_i, B_i \in \mathcal{B}(\mathcal{H}).$$

Because the detailed physically plausible conditions on the systems, which correspond to these properties are not known, it is much more convenient to adopt an axiomatic point of view which is based mainly on the simplicity and the succes of physical applications. Accordingly [2,9,13,14,30] it is convenient to suppose that the time evolutions Φ_t for open systems are not very different from the time evolutions for closed systems. The simplest dynamics Φ_t which introduces a preferred direction in time, characteristic for dissipative processes, is that in which the group condition is replaced by the semigroup condition [9-12,31]

$$\Phi_{t+s} = \Phi_t \Phi_s, \quad t, s \geq 0.$$

The duality condition

$$\text{Tr}(\Phi_t(\rho) A) = \text{Tr}(\rho \tilde{\Phi}_t(A)) \tag{2.1}$$

defines $\tilde{\Phi}_t$, the dual of Φ_t acting on $\mathcal{B}(\mathcal{H})$. Then the conditions

$$\mathrm{Tr}\Phi_t(\rho) = 1$$

and

$$\tilde{\Phi}_t(I) = I \tag{2.2}$$

are equivalent. Also the conditions

$$\tilde{\Phi}_t(A) \rightarrow A \tag{2.3}$$

ultraweakly when $t \rightarrow 0$ and

$$\Phi_t(\rho) \rightarrow \rho$$

in the trace norm when $t \rightarrow 0$, are equivalent. For the semigroups with the properties (2.2), (2.3) and

$$A \geq 0 \rightarrow \tilde{\Phi}_t(A) \geq 0,$$

it is well known that there exists a (generally bounded) mapping $\tilde{L}$, the generator of $\tilde{\Phi}_t$. $\tilde{\Phi}_t$ is uniquely determined by $\tilde{L}$. The dual generator of the dual semigroup Φ_t is denoted by L :

$$\mathrm{Tr}(L(\rho)A) = \mathrm{Tr}(\rho\tilde{L}(A)).$$

The evolution equations by which L and $\tilde{L}$ determine uniquely Φ_t and $\tilde{\Phi}_t$, respectively, are given in the Schrödinger and Heisenberg picture as

$$\frac{d\Phi_t(\rho)}{dt} = L(\Phi_t(\rho)) \tag{2.4}$$

and

$$\frac{d\tilde{\Phi}_t(A)}{dt} = \tilde{L}(\tilde{\Phi}_t(A)). \tag{2.5}$$

These equations replace in the case of open systems the von Neumann-Liouville equations

$$\frac{dU_t(\rho)}{dt} = -\frac{i}{\hbar}[H, U_t(\rho)]$$

and

$$\frac{d\tilde{U}_t(A)}{dt} = \frac{i}{\hbar}[H, \tilde{U}_t(A)].$$

For any applications Eqs. (2.4) and (2.5) are only useful if the detailed structure of the generator $L(\tilde{L})$ is known and can be related to the concrete properties of the open systems described by such equations.

Such a structural theorem was obtained by Lindblad [14] for the class of dynamical semigroups $\tilde{\Phi}_t$ which are completely positive and norm continuous. For such semigroups the generator $\tilde{L}$ is bounded. In many applications the generator is unbounded.

A bounded mapping $\tilde{L} : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ which satisfies $\tilde{L}(I) = 0$, $\tilde{L}(A^+) = \tilde{L}(A)^+$ and

$$\tilde{L}(A^+A) - \tilde{L}(A^+)A - A^+\tilde{L}(A) \geq 0$$

is called dissipative. The 2-positivity property of the completely positive mapping $\tilde{\Phi}_t$:

$$\tilde{\Phi}_t(A^+A) \geq \tilde{\Phi}_t(A^+)\tilde{\Phi}_t(A), \quad (2.6)$$

with equality at $t = 0$, implies that $\tilde{L}$ is dissipative. Lindblad [14] has shown that conversely, the dissipativity of $\tilde{L}$ implies that $\tilde{\Phi}_t$ is 2-positive. $\tilde{L}$ is called completely dissipative if all trivial extensions of $\tilde{L}$ are dissipative. Lindblad has also shown that there exists a one-to-one correspondence between the completely positive norm continuous semigroups $\tilde{\Phi}_t$ and completely dissipative generators $\tilde{L}$. The structural theorem of Lindblad gives the most general form of a completely dissipative mapping $\tilde{L}$ [14]:

Theorem: $\tilde{L}$ is completely dissipative and ultraweakly continuous if and only if it is of the form

$$\tilde{L}(A) = \frac{i}{\hbar}[H, A] + \frac{1}{2\hbar} \sum_j (V_j^+[A, V_j] + [V_j^+, A]V_j), \quad (2.7)$$

where $V_j, \sum_j V_j^+ V_j \in \mathcal{B}(\mathcal{H})$, $H \in \mathcal{B}(\mathcal{H})_{\text{s.a.}}$.

The dual generator on the state space (Schrödinger picture) is of the form

$$L(\rho) = -\frac{i}{\hbar}[H, \rho] + \frac{1}{2\hbar} \sum_j ([V_j \rho, V_j^+] + [V_j, \rho V_j^+]). \quad (2.8)$$

Eqs. (2.4) and (2.8) give an explicit form for the most general time-homogeneous quantum mechanical Markovian master equation with a bounded Liouville operator:

$$\frac{d\Phi_t(\rho)}{dt} = L(\Phi_t(\rho)) = -\frac{i}{\hbar}[H, \Phi_t(\rho)] + \frac{1}{2\hbar} \sum_j ([V_j \Phi_t(\rho), V_j^+] + [V_j, \Phi_t(\rho) V_j^+]).$$

Talkner [32] has shown that the assumption of a semigroup dynamics is only applicable in the limit of weak coupling of the subsystem with its environment, i.e. for long relaxation times.

We should like to mention that all Markovian master equations found in the literature are of this form after some rearrangement of terms, even for unbounded generators.

It is also an empirical fact for many physically interesting situations that the time evolutions Φ_t drive the system towards a unique final state $\rho(\infty) = \lim_{t \rightarrow \infty} \Phi_t(\rho(0))$ for all $\rho(0) \in \mathcal{D}(\mathcal{H})$.

From the 2-positivity property (2.6) of $\tilde{\Phi}_t$ it follows that

$$\tilde{\Phi}(t)(\sum_j V_j^* V_j) \geq \sum_j \tilde{\Phi}_t(V_j^*) \tilde{\Phi}_t(V_j)$$

or, by duality [17],

$$\text{Tr}(\tilde{\Phi}_t(\rho) \sum_j V_j^* V_j) \geq \sum_j \text{Tr}(\tilde{\Phi}_t(\rho) V_j^*) \text{Tr}(\tilde{\Phi}_t(\rho) V_j). \quad (2.9)$$

The evolution equations of Lindblad are operator equations. The problem of finding their solutions is in general rather difficult. In cases when the equations are exactly solvable, these solutions give complete information about the studied problem and permit the calculation of expectation values of the observables at any moment.

3. The one-dimensional damped quantum harmonic oscillator

In this Section the case of the damped quantum harmonic oscillator is considered in the spirit of the ideas presented in the previous Section. The basic assumption is that the general form (2.8) of a bounded mapping L given by the Lindblad theorem [14] is also valid for an unbounded completely dissipative mapping L :

$$L(\rho) = -\frac{i}{\hbar}[H, \rho] + \frac{1}{2\hbar} \sum_j ([V_j \rho, V_j^+] + [V_j, \rho V_j^+]). \quad (3.1)$$

This assumption gives one of the simplest way to construct an appropriate model for a quantum dissipative system. Another simple condition imposed to the operators H, V_j, V_j^+ is that they are functions of the basic observables q and p of the one-dimensional quantum mechanical system (with $[q, p] = i\hbar I$, where I is the identity operator on $\mathcal{H}$) of such kind that the obtained model is exactly solvable. A precise version for this last condition is

that linear spaces spanned by the first degree (respectively second degree) noncommutative polynomials in p and q are invariant under the action of the completely dissipative mapping L . This condition implies [16] that V_j are at most first degree polynomials in p and q and H is at most a second degree polynomial in p and q .

Because in the linear space of the first degree polynomials in p and q the operators p and q give a basis, there exist only two $\mathbf{C}$ -linear independent operators V_1, V_2 which can be written in the form

$$V_i = a_i p + b_i q, \quad i = 1, 2,$$

with $a_i, b_i = 1, 2$ complex numbers [16]. The constant term is omitted because its contribution to the generator L is equivalent to terms in H linear in p and q which for simplicity are assumed to be zero. Then H is chosen of the form

$$H = H_0 + \frac{\mu}{2}(pq + qp), \quad H_0 = \frac{1}{2m}p^2 + \frac{m\omega^2}{2}q^2. \quad (3.2)$$

With these choices the Markovian master equation can be written:

$$\begin{aligned} \frac{d\rho}{dt} = & -\frac{i}{\hbar}[H_0, \rho] - \frac{i}{2\hbar}(\lambda + \mu)[q, \rho p + p\rho] + \frac{i}{2\hbar}(\lambda - \mu)[p, \rho q + q\rho] \\ & - \frac{D_{pp}}{\hbar^2}[q, [q, \rho]] - \frac{D_{qq}}{\hbar^2}[p, [p, \rho]] + \frac{D_{pq}}{\hbar^2}([q, [p, \rho]] + [p, [q, \rho]]). \end{aligned} \quad (3.3)$$

Here we used the notations:

$$D_{qq} = \frac{\hbar}{2} \sum_{j=1,2} |a_j|^2, D_{pp} = \frac{\hbar}{2} \sum_{j=1,2} |b_j|^2, D_{pq} = D_{qp} = -\frac{\hbar}{2} \text{Re} \sum_{j=1,2} a_j^* b_j, \lambda = -\text{Im} \sum_{j=1,2} a_j^* b_j,$$

where D_{pp}, D_{qq} and D_{pq} are the diffusion coefficients and λ the friction constant. They satisfy the following fundamental constraints [17]:

$$\begin{aligned} i) \quad & D_{pp} > 0, \\ ii) \quad & D_{qq} > 0, \\ iii) \quad & D_{pp}D_{qq} - D_{pq}^2 \geq \lambda^2 \hbar^2 / 4. \end{aligned} \quad (3.4)$$

Indeed, inequalities $i)$ and $ii)$ follow directly from the definitions of the coefficients and $iii)$ from the Schwartz inequality. The equality $\mu = \lambda$ is a necessary and sufficient condition

for L to be translation invariant [16]: $[p, L(\rho)] = L([p, \rho])$. In the following general values for λ and μ will be considered as in [16].

Introducing the annihilation and creation operators

$$a = \frac{1}{\sqrt{2\hbar}}(\sqrt{m\omega}q + \frac{i}{\sqrt{m\omega}}p), \quad a^+ = \frac{1}{\sqrt{2\hbar}}(\sqrt{m\omega}q - \frac{i}{\sqrt{m\omega}}p), \quad (3.5)$$

obeying the commutation relation $[a, a^+] = 1$, we have

$$H_0 = \hbar\omega(a^+a + \frac{1}{2}) \quad (3.6)$$

and the master equation has the form

$$\begin{aligned} \frac{d\rho}{dt} = & \frac{1}{2}(D_1 - \mu)(\rho a^+ a^+ - a^+ \rho a^+) + \frac{1}{2}(D_1 + \mu)(a^+ a^+ \rho - a^+ \rho a^+) \\ & + \frac{1}{2}(D_2 - \lambda - i\omega)(a^+ \rho a - \rho a a^+) + \frac{1}{2}(D_2 + \lambda + i\omega)(a \rho a^+ - a^+ a \rho) + h.c., \end{aligned} \quad (3.7)$$

where

$$D_1 = \frac{1}{\hbar}(m\omega D_{qq} - \frac{D_{pp}}{m\omega} + 2iD_{pq}), \quad D_2 = \frac{1}{\hbar}(m\omega D_{qq} + \frac{D_{pp}}{m\omega}). \quad (3.8)$$

In the literature, master equations of the type (3.3) or (3.7) are encountered in models for the description of different physical phenomena in quantum optics, in treatments of the damping of collective modes in deep inelastic collisions of heavy ions or in the quantum mechanical description of the dissipation for the one-dimensional harmonic oscillator. In the following we show that some of these master equations are particular cases of the Lindblad equation.

1) The master equation of Dekker for the damped quantum harmonic oscillator [5,33-36] supplemented with the fundamental constraints (3.4) obtained in [36] from the condition that the time evolution of this master equation does not violate the generalized uncertainty principle [37] at any time, is a particular case of the Lindblad master equation (3.7) when $\mu = \lambda$. If the constraints (3.4) are not fulfilled in the case $\mu = \lambda$, then the uncertainty principle is violated.

2) The quantum master equation considered in [38,39] by Hofmann et al. for treating the charge equilibration process as a collective high frequency mode is a particular case of the Lindblad master equation (3.3) if $\lambda = \gamma(\omega)/2m = \mu$, $D_{qq} = D_{pq} = 0$, $D_{pp} = \gamma(\omega)T^*(\omega)$, but the fundamental constraints (3.4) are not satisfied.

3) For the quantum master equation considered in [40] for the description of heavy ion collisions we have $\lambda = \mu = \gamma/2, D_{pp} = D, D_{qq} = 0, D_{pq} = D_{qp} = -d/2$ and consequently the fundamental constraints are not fulfilled.

4) In [41], Spina and Weidenmüller considered two kinds of master equations for describing the damping of collective modes in deep inelastic collisions of heavy ions.

(i) The first equation can be obtained from Eq. (3.3) by replacing H_0 by

$H_0 - Am\omega q^2/2 + f(t)q$ and setting $\lambda = \mu = \Gamma/2, D_{pp} = D/2, D_{qq} = 0$ and $D_{pq} = D_{qp} = B/2$. Then the constraints (3.4) are not satisfied.

(ii) The second master equation is obtained from (3.3) by setting $H_0 - Am\omega q^2/2 - Ap^2/(2m\omega) + f(t)q$ instead of H_0 and $\Gamma_R = \Gamma_p = \lambda, \mu = 0, D_{pp} = D_p/2, D_{qq} = D_R/2, D_{pq} = 0$. Then last condition (3.4) is satisfied for all values of the parameters.

5) The master equation for the density operator of the electromagnetic field mode coupled to a squeezed bath [42,43] can be obtained from the master equation (3.7) if we set $\mu = 0, \lambda = \gamma, D_1 = 2\gamma M, D_2 = \gamma(2N + 1)$ (in [42]: $\gamma/2$ instead of γ and $\bar{N}$ instead of N).

6) The master equation for the density operator of a harmonic oscillator coupled to an environment of harmonic oscillators considered in [44-47] is a particular case of the master equation (3.7) if we set $\lambda = \mu = \gamma, D_{qq} = D_{pq} = 0, D_{pp} = 2\gamma(\bar{n} + 1/2)m\omega\hbar$ and the fundamental constraints (3.4) are not fulfilled.

7) The master equation written in [48] for different models of correlated- emission lasers can also be obtained from the master equation (3.7) by setting

$$D_1 + \mu = 2\Lambda_4, D_1 - \mu = 2\Lambda_3, D_2 + \lambda + i\omega = 2\Lambda_2, D_2 - \lambda - i\omega = 2\Lambda_1.$$

8) Two master equations were introduced by Jang and Yannouleas in [21,49], where the nuclear dissipative process is described as the damping of a collective degree of freedom coupled to a bosonic reservoir at finite temperature. The resulting RPA master equation within the observed collective subspace was derived in a purely dynamical way.

(i) The master equation written in [49] in the resonant approximation (rotating-wave approximation) can be obtained as a particular case of the Lindblad master equation (3.7). For this one has to set

$$D_{pp} = m^2\omega^2 D_{qq}, D_{pq} = \mu = 0, \frac{4m\omega D_{qq}}{\hbar} = (2\langle n \rangle + 1)\Gamma, \lambda = \frac{\Gamma}{2},$$

where $\langle n \rangle$ is the average number of the RPA collective phonons at thermal equilibrium and Γ the decay constant. The fundamental constraints (3.4) are fulfilled in this case.

(ii) The master equation derived in [21] in order to extend the calculations carried out in [49] with the before-mentioned master equation, can also be obtained as a particular case of the master equation (3.7) by setting

$$D_{qq} = D_{pq} = 0, \quad D_{pp} = \frac{\hbar m \omega}{2} (2 \langle n \rangle + 1) \Gamma, \quad \mu = \lambda = \frac{\Gamma}{2}$$

or $D_2 = -D_1 = (2 \langle n \rangle + 1) \Gamma / 2$. In this case the fundamental constraints (3.4) are not fulfilled.

As a conclusion, we can say that on various occasions the master equation for $\lambda = \mu$ has been tested in the sense that it does not violate the uncertainty principle in the final steady state, but a possible transient violation of the uncertainty principle has been noted [5,36]. The above mentioned models can be divided into three types. The first type are models which fulfill the constraints (3.4). The corresponding master equations belong to the category of equations described by the Lindblad theory. These models obey the uncertainty principle. The second type are models with $\lambda = \mu$ which do not fulfill the constraints (3.4) (for example model 2). According to [36] these models show a dynamical violation of the uncertainty principle. The third type are models with $\lambda \neq \mu$ which also do not fulfill the constraints (3.4). These models and those of the second type do not belong to the category of master equations treated in the Lindblad theory. For the models of the third type we can not prove that they do not conserve the uncertainty principle in time.

3.1 The master equation in the Heisenberg picture

The following notations will be used:

$$\begin{aligned} \sigma_q(t) &= \text{Tr}(\rho(t)q), \\ \sigma_p(t) &= \text{Tr}(\rho(t)p), \\ \sigma_{qq} &= \text{Tr}(\rho(t)q^2) - \sigma_q^2(t), \\ \sigma_{pp} &= \text{Tr}(\rho(t)p^2) - \sigma_p^2(t), \\ \sigma_{pq}(t) &= \text{Tr}(\rho(t) \frac{pq + qp}{2}) - \sigma_p(t)\sigma_q(t). \end{aligned} \tag{3.9}$$

In the Heisenberg picture the master equation has the following symmetric form:

$$\begin{aligned}
\frac{d\tilde{\Phi}_t(A)}{dt} &= \tilde{L}(\tilde{\Phi}_t(A)) = \frac{i}{\hbar}[H_0, \tilde{\Phi}_t(A)] - \frac{i}{2\hbar}(\lambda + \mu)([\tilde{\Phi}_t(A), q]p + p[\tilde{\Phi}_t(A), q]) \\
&+ \frac{i}{2\hbar}(\lambda - \mu)(q[\tilde{\Phi}_t(A), p] + [\tilde{\Phi}_t(A), p]q) - \frac{D_{pp}}{\hbar^2}[q, [q, \tilde{\Phi}_t(A)]] \\
&- \frac{D_{qq}}{\hbar^2}[p, [p, \tilde{\Phi}_t(A)]] + \frac{D_{pq}}{\hbar^2}([p, [q, \tilde{\Phi}_t(A)]] + [q, [p, \tilde{\Phi}_t(A)]]). \tag{3.10}
\end{aligned}$$

Denoting by A a selfadjoint operator we have

$$\sigma_A(t) = \text{Tr}(\rho(t)A), \sigma_{AA}(t) = \text{Tr}(\rho(t)A^2) - \sigma_A^2(t).$$

It follows that

$$\frac{d\sigma_A(t)}{dt} = \text{Tr}(L(\rho(t))A) = \text{Tr}(\rho(t)\tilde{L}(A)) \tag{3.11}$$

and

$$\frac{d\sigma_{AA}(t)}{dt} = \text{Tr}(L(\rho(t)A^2) - 2\frac{d\sigma_A(t)}{dt}\sigma_A(t)) = \text{Tr}(\rho(t)\tilde{L}(A^2)) - 2\sigma_A(t)\text{Tr}((\rho(t)\tilde{L}(A))). \tag{3.12}$$

An important consequence of the precise version of solvability condition formulated at the beginning of the present Section is the fact that when A is put equal to p or q in (3.11) and (3.12), then $d\sigma_p(t)/dt$ and $d\sigma_q(t)/dt$ are functions only of $\sigma_p(t)$ and $\sigma_q(t)$ and $d\sigma_{pp}(t)/dt$, $d\sigma_{qq}(t)/dt$ and $d\sigma_{pq}(t)/dt$ are functions only of $\sigma_{pp}(t)$, $\sigma_{qq}(t)$ and $\sigma_{pq}(t)$. This fact allows an immediate determination of $\sigma_p(t)$, $\sigma_q(t)$, $\sigma_{pp}(t)$, $\sigma_{qq}(t)$ and $\sigma_{pq}(t)$ as functions of time. The results are the following:

$$\begin{aligned}
\frac{d\sigma_q(t)}{dt} &= -(\lambda - \mu)\sigma_q(t) + \frac{1}{m}\sigma_p(t), \\
\frac{d\sigma_p(t)}{dt} &= -m\omega^2\sigma_q(t) - (\lambda + \mu)\sigma_p(t) \tag{3.13}
\end{aligned}$$

and

$$\begin{aligned}
\frac{d\sigma_{qq}(t)}{dt} &= -2(\lambda - \mu)\sigma_{qq}(t) + \frac{2}{m}\sigma_{pq}(t) + 2D_{qq}, \\
\frac{d\sigma_{pp}(t)}{dt} &= -2(\lambda + \mu)\sigma_{pp}(t) - 2m\omega^2\sigma_{pq}(t) + 2D_{pp}, \\
\frac{d\sigma_{pq}(t)}{dt} &= -m\omega^2\sigma_{qq}(t) + \frac{1}{m}\sigma_{pp}(t) - 2\lambda\sigma_{pq}(t) + 2D_{pq}. \tag{3.14}
\end{aligned}$$

All equations considered in various papers in connection with damping of collective modes in deep inelastic collisions are obtained as particular cases of Eqs. (3.13) and (3.14), as we already mentioned before.

The integration of Eqs. (3.13) is straightforward. There are two cases: *a)* $\mu > \omega$ (overdamped) and *b)* $\mu < \omega$ (underdamped).

In the case *a)* with the notation $\nu^2 = \mu^2 - \omega^2$, we obtain [17]:

$$\begin{aligned}\sigma_q(t) &= e^{-\lambda t} \left((\cosh \nu t + \frac{\mu}{\nu} \sinh \nu t) \sigma_q(0) + \frac{1}{m\nu} \sinh \nu t \sigma_p(0) \right), \\ \sigma_p(t) &= e^{-\lambda t} \left(-\frac{m\omega^2}{\nu} \sinh \nu t \sigma_q(0) + (\cosh \nu t - \frac{\mu}{\nu} \sinh \nu t) \sigma_p(0) \right).\end{aligned}\quad (3.15)$$

If $\lambda > \nu$, then $\sigma_q(\infty) = \sigma_p(\infty) = 0$. If $\lambda < \nu$, then $\sigma_q(\infty) = \sigma_p(\infty) = \infty$.

In the case *b)* with the notation $\Omega^2 = \omega^2 - \mu^2$, we obtain [17]:

$$\begin{aligned}\sigma_q(t) &= e^{-\lambda t} \left((\cos \Omega t + \frac{\mu}{\Omega} \sin \Omega t) \sigma_q(0) + \frac{1}{m\Omega} \sin \Omega t \sigma_p(0) \right), \\ \sigma_p(t) &= e^{-\lambda t} \left(-\frac{m\omega^2}{\Omega} \sin \Omega t \sigma_q(0) + (\cos \Omega t - \frac{\mu}{\Omega} \sin \Omega t) \sigma_p(0) \right)\end{aligned}\quad (3.16)$$

and $\sigma_q(\infty) = \sigma_p(\infty) = 0$. In order to integrate Eqs. (3.14), it is convenient to consider the vector

$$X(t) = \begin{pmatrix} m\omega\sigma_{qq}(t) \\ \sigma_{pp}(t)/m\omega \\ \sigma_{pq}(t) \end{pmatrix}.$$

Then the system of equations (3.14) can be written in the form

$$\frac{dX(t)}{dt} = RX(t) + D,$$

where R is the following 3×3 matrix

$$R = \begin{pmatrix} -2(\lambda - \mu) & 0 & 2\omega \\ 0 & -2(\lambda + \mu) & -2\omega \\ -\omega & \omega & -2\lambda \end{pmatrix}$$

and D the following vector

$$D = \begin{pmatrix} 2m\omega D_{qq} \\ 2D_{pp}/m\omega \\ 2D_{pq} \end{pmatrix}.$$

Then there exists a matrix T with the property $T^2 = I$ where I is the identity matrix and a diagonal matrix K such that $R = TKT$. From this it follows that

$$X(t) = (Te^{Kt}T)X(0) + T(e^{Kt} - I)K^{-1}TD. \quad (3.17)$$

An interesting observation is that the time dependence of the variances $\sigma_{qq}(t)$, $\sigma_{pp}(t)$, $\sigma_{pq}(t)$ decomposes in a classical part given by $Te^{Kt}TX(0)$ and a quantum part given by $T(e^{Kt} - I)K^{-1}TD$. Exactly this quantum part governs the asymptotic behaviour of the variances when $t \rightarrow \infty$ [17].

In the overdamped case ($\mu > \omega$) the matrices T and K are given by

$$T = \frac{1}{2\nu} \begin{pmatrix} \mu + \nu & \mu - \nu & 2\omega \\ \mu - \nu & \mu + \nu & 2\omega \\ -\omega & -\omega & -2\mu \end{pmatrix}$$

and

$$K = \begin{pmatrix} -2(\lambda - \nu) & 0 & 0 \\ 0 & -2(\lambda + \nu) & 0 \\ 0 & 0 & -2\lambda \end{pmatrix}$$

with $\nu^2 = \mu^2 - \omega^2$.

In the underdamped case ($\mu < \omega$) the matrices T and K are given by

$$T = \frac{1}{2i\Omega} \begin{pmatrix} \mu + i\Omega & \mu - i\Omega & 2\omega \\ \mu - i\Omega & \mu + i\Omega & 2\omega \\ -\omega & -\omega & -2\mu \end{pmatrix}$$

and

$$K = \begin{pmatrix} -2(\lambda - i\Omega) & 0 & 0 \\ 0 & -2(\lambda + i\Omega) & 0 \\ 0 & 0 & -2\lambda \end{pmatrix}$$

with $\Omega^2 = \omega^2 - \mu^2$. From (3.17) it follows that

$$X(\infty) = -(TK^{-1}T)D = -R^{-1}D \quad (3.18)$$

(in the overdamped case the restriction $\lambda > \nu$ is necessary). Then Eq.(3.17) can be written in the form

$$X(t) = (Te^{Kt}T)(X(0) - X(\infty)) + X(\infty). \quad (3.19)$$

Also

$$\frac{dX(t)}{dt} = (TKe^{Kt}T)(X(0) - X(\infty)) = R(X(t) - X(\infty)).$$

The formula (3.18) is remarkable because it gives a very simple connection between the asymptotic values ($t \rightarrow \infty$) of $\sigma_{qq}(t), \sigma_{pp}(t), \sigma_{pq}(t)$ and the diffusion coefficients D_{qq}, D_{pp}, D_{pq} . As an immediate consequence of (3.18) this connection is *the same* for both cases, underdamped and overdamped, and has the following explicit form:

$$\begin{aligned}\sigma_{qq}(\infty) &= \frac{1}{2(m\omega)^2\lambda(\lambda^2 + \omega^2 - \mu^2)}((m\omega)^2(2\lambda(\lambda + \mu) + \omega^2)D_{qq} \\ &\quad + \omega^2 D_{pp} + 2m\omega^2(\lambda + \mu)D_{pq}), \\ \sigma_{pp}(\infty) &= \frac{1}{2\lambda(\lambda^2 + \omega^2 - \mu^2)}((m\omega)^2\omega^2 D_{qq} + (2\lambda(\lambda - \mu) + \omega^2)D_{pp} - 2m\omega^2(\lambda - \mu)D_{pq}), \quad (3.20) \\ \sigma_{pq}(\infty) &= \frac{1}{2m\lambda(\lambda^2 + \omega^2 - \mu^2)}(-(\lambda + \mu)(m\omega)^2 D_{qq} + (\lambda - \mu)D_{pp} + 2m(\lambda^2 - \mu^2)D_{pq}).\end{aligned}$$

These relations show that the asymptotic values $\sigma_{qq}(\infty), \sigma_{pp}(\infty), \sigma_{pq}(\infty)$ do not depend on the initial values $\sigma_{qq}(0), \sigma_{pp}(0), \sigma_{pq}(0)$. In other words,

$$R^{-1} = \frac{-1}{4\lambda(\lambda^2 + \omega^2 - \mu^2)} \begin{pmatrix} 2\lambda(\lambda + \mu) + \omega^2 & \omega^2 & 2\omega(\lambda + \mu) \\ \omega^2 & 2\lambda(\lambda - \mu) + \omega^2 & -2\omega(\lambda - \mu) \\ -(\lambda + \mu)\omega & (\lambda - \mu)\omega & 2(\lambda^2 - \mu^2) \end{pmatrix}.$$

Conversely, if the relations $D = -RX(\infty)$ are considered, then

$$\begin{aligned}D_{qq} &= (\lambda - \mu)\sigma_{qq}(\infty) - \frac{1}{m}\sigma_{pq}(\infty), \\ D_{pp} &= (\lambda + \mu)\sigma_{pp}(\infty) + m\omega^2\sigma_{pq}(\infty), \\ D_{pq} &= \frac{1}{2}(m\omega^2\sigma_{qq}(\infty) - \frac{1}{m}\sigma_{pp}(\infty) + 2\lambda\sigma_{pq}(\infty)).\end{aligned} \quad (3.21)$$

Hence, from (3.4) fundamental constraints on $\sigma_{qq}(\infty), \sigma_{pp}(\infty)$ and $\sigma_{pq}(\infty)$ follow:

$$\begin{aligned}D_{qq} &= (\lambda - \mu)\sigma_{qq}(\infty) - \frac{1}{m}\sigma_{pq}(\infty) > 0, \\ D_{pp} &= (\lambda + \mu)\sigma_{pp}(\infty) + m\omega^2\sigma_{pq}(\infty) > 0, \\ D_{qq}D_{pp} - D_{pq}^2 &= (\lambda^2 - \mu^2)\sigma_{qq}(\infty)\sigma_{pp}(\infty) - \omega^2\sigma_{pq}^2(\infty) + (\lambda - \mu)m\omega^2\sigma_{qq}(\infty)\sigma_{pq}(\infty) \\ &\quad - \frac{(\lambda + \mu)}{m}\sigma_{pp}(\infty)\sigma_{pq}(\infty) - \frac{1}{4}(m\omega^2)^2\sigma_{qq}^2(\infty) - \frac{1}{4m^2}\sigma_{pp}^2(\infty) - \lambda^2\sigma_{pq}^2(\infty) + \frac{1}{2}\omega^2\sigma_{qq}(\infty)\sigma_{pp}(\infty) \\ &\quad - m\omega^2\lambda\sigma_{qq}(\infty)\sigma_{pq}(\infty) + \frac{\lambda}{m}\sigma_{pp}(\infty)\sigma_{pq}(\infty) \geq \frac{\lambda^2\hbar^2}{4}.\end{aligned} \quad (3.22)$$

The constraint (3.22) can be put in a more clear form:

$$4(\lambda^2 + \omega^2 - \mu^2)(\sigma_{qq}(\infty)\sigma_{pp}(\infty) - \sigma_{pq}(\infty)^2) - (m\omega^2\sigma_{qq}(\infty) + \frac{1}{m}\sigma_{pp}(\infty) + 2\mu\sigma_{pq}^2(\infty)) \geq \hbar^2\lambda^2. \quad (3.23)$$

If $\mu < \omega$ (the underdamped case), then $\lambda^2 + \omega^2 - \mu^2 > \lambda^2$. If $\mu > \omega$ (the overdamped case), then $0 \leq \lambda^2 + \omega^2 - \mu^2 < \lambda^2$ ($\lambda > \nu$) and the constraint (3.23) is more strong than the uncertainty inequality $\sigma_{qq}(\infty)\sigma_{pp}(\infty) - \sigma_{pq}^2(\infty) \geq \hbar^2/4$. By using the fact that the linear positive mapping $\mathcal{B}(\mathcal{H}) \rightarrow \mathbf{C}$ defined by $A \rightarrow \text{Tr}(\rho A)$ is completely positive (hence 2-positive), in [17] the following inequality was obtained:

$$D_{qq}\sigma_{pp}(t) + D_{pp}\sigma_{qq}(t) - 2D_{pq}\sigma_{pq}(t) \geq \frac{\hbar^2\lambda}{2}.$$

We found this inequality, which must be valid for all values of $t \in (0, \infty)$, as the sufficient condition that the generalized uncertainty inequality [37]

$$\sigma_{qq}(t)\sigma_{pp}(t) - \sigma_{pq}^2(t) \geq \frac{\hbar^2}{4}$$

is fulfilled at any moment t , if the initial values $\sigma_{qq}(0), \sigma_{pp}(0)$ and $\sigma_{pq}(0)$ for $t = 0$ satisfy this inequality. A restriction connecting the initial values $\sigma_{qq}(0), \sigma_{pp}(0), \sigma_{pq}(0)$ with the asymptotic values $\sigma_{pp}(\infty), \sigma_{qq}(\infty), \sigma_{pq}(\infty)$ is also obtained if the values D_{qq}, D_{pp} and D_{pq} are expressed by Eq. (3.21) in terms of $\sigma_{pp}(\infty), \sigma_{qq}(\infty), \sigma_{pq}(\infty)$:

$$D_{qq}\sigma_{pp}(0) + D_{pp}\sigma_{qq}(0) - 2D_{pq}\sigma_{pq}(0) \geq \frac{\hbar^2\lambda}{2}.$$

More explicitly

$$\begin{aligned} & \lambda(\sigma_{qq}(\infty)\sigma_{pp}(0) + \sigma_{pp}(\infty)\sigma_{qq}(0) - 2\sigma_{pq}(\infty)\sigma_{pq}(0)) - \mu(\sigma_{qq}(\infty)\sigma_{pp}(0) - \sigma_{pp}(\infty)\sigma_{qq}(0)) \\ & - \frac{1}{m}(\sigma_{pq}(\infty)\sigma_{pp}(0) - \sigma_{pp}(\infty)\sigma_{pq}(0)) + m\omega^2(\sigma_{pq}(\infty)\sigma_{qq}(0) - \sigma_{qq}(\infty)\sigma_{pq}(0)) \geq \frac{\hbar^2\lambda}{2}. \end{aligned} \quad (3.24)$$

If the asymptotic state is a Gibbs state

$$\rho_G(\infty) = e^{-H_0/kT} / \text{Tr}(e^{-H_0/kT}),$$

then

$$\sigma_{qq}(\infty) = \frac{\hbar}{2m\omega} \coth \frac{\hbar\omega}{2kT}, \quad \sigma_{pp}(\infty) = \frac{\hbar m\omega}{2} \coth \frac{\hbar\omega}{2kT}, \quad \sigma_{pq}(\infty) = 0 \quad (3.25)$$

and

$$D_{pp} = \frac{\lambda + \mu}{2} \hbar m\omega \coth \frac{\hbar\omega}{2kT}, \quad D_{qq} = \frac{\lambda - \mu}{2} \frac{\hbar}{m\omega} \coth \frac{\hbar\omega}{2kT}, \quad D_{pq} = 0 \quad (3.26)$$

and the fundamental constraints (3.4) are satisfied only if $\lambda > \mu$ and [16]:

$$(\lambda^2 - \mu^2) \left(\coth \frac{\hbar\omega}{2kT} \right)^2 \geq \lambda^2.$$

If the initial state is the ground state of the harmonic oscillator, then

$$\sigma_{qq}(0) = \frac{\hbar}{2m\omega}, \quad \sigma_{pp}(0) = \frac{m\hbar\omega}{2}, \quad \sigma_{pq}(0) = 0.$$

Then (3.24) becomes

$$\lambda \left(\sigma_{qq}(\infty) m\omega + \frac{\sigma_{pp}(\infty)}{m\omega} \right) - \mu \left(\sigma_{qq}(\infty) m\omega - \frac{\sigma_{pp}(\infty)}{m\omega} \right) \geq \hbar \lambda.$$

For example, in the case (3.25), this implies $\coth(\hbar\omega/2kT) \geq 1$ which is always valid.

Now, the explicit time dependence of $\sigma_{qq}(t), \sigma_{pp}(t), \sigma_{pq}(t)$ will be given for both under- and overdamped cases. From Eq. (3.19) it follows that in order to obtain the explicit time dependence it is necessary to obtain the matrix elements of $Te^{Kt}T$. In the overdamped case ($\mu > \omega, \nu^2 = \mu^2 - \omega^2$) we have

$$Te^{Kt}T = \frac{e^{-2\lambda t}}{2\nu^2} \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix},$$

with

$$\begin{aligned} a_{11} &= (\mu^2 + \nu^2) \cosh 2\nu t + 2\mu\nu \sinh 2\nu t - \omega^2, \\ a_{12} &= (\mu^2 - \nu^2) \cosh 2\nu t - \omega^2, \\ a_{13} &= 2\omega(\mu \cosh 2\nu t + \nu \sinh 2\nu t - \mu), \\ a_{21} &= (\mu^2 - \nu^2) \cosh 2\nu t - \omega^2, \\ a_{22} &= (\mu^2 + \nu^2) \cosh 2\nu t - 2\mu\nu \sinh 2\nu t - \omega^2, \\ a_{23} &= 2\omega(\mu \cosh 2\nu t - \nu \sinh 2\nu t - \mu), \\ a_{31} &= -\omega(\mu \cosh 2\nu t + \nu \sinh 2\nu t - \mu), \\ a_{32} &= -\omega(\mu \cosh 2\nu t - \nu \sinh 2\nu t - \mu), \end{aligned} \quad (3.27)$$

$$a_{33} = -2(\omega^2 \cosh 2\nu t - \mu^2).$$

In the underdamped case ($\mu < \omega$, $\Omega^2 = \omega^2 - \mu^2$) we have

$$Te^{Kt}T = -\frac{e^{-2\lambda t}}{2\Omega^2} \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{pmatrix}$$

with

$$\begin{aligned} b_{11} &= (\mu^2 - \Omega^2) \cos 2\Omega t - 2\mu\Omega \sin 2\Omega t - \omega^2, \\ b_{12} &= (\mu^2 + \Omega^2) \cos 2\Omega t - \omega^2, \\ b_{13} &= 2\omega(\mu \cos 2\Omega t - \Omega \sin 2\Omega t - \mu), \\ b_{21} &= (\mu^2 + \Omega^2) \cos 2\Omega t - \omega^2, \\ b_{22} &= (\mu^2 - \Omega^2) \cos 2\Omega t + 2\mu\Omega \sin 2\Omega t - \omega^2, \\ b_{23} &= 2\omega(\mu \cos 2\Omega t + \Omega \sin 2\Omega t - \mu), \\ b_{31} &= -\omega(\mu \cos 2\Omega t - \Omega \sin 2\Omega t - \mu), \\ b_{32} &= -\omega(\mu \cos 2\Omega t + \Omega \sin 2\Omega t - \mu), \\ b_{33} &= -2(\omega^2 \cos 2\Omega t - \mu^2). \end{aligned} \tag{3.28}$$

As was shown by Lindblad in [16] the equations of motion (3.10) written for Weyl operators ($A = W(\xi, \eta) = e^{(i/\hbar)(\eta q - \xi p)}$) can be integrated in a very simple and elegant way. This fact is important from both points of view: practical and theoretical. From practical point of view, it gives in a new way explicit formulas for $\sigma_{qq}(t)$, $\sigma_{pp}(t)$, $\sigma_{pq}(t)$ and moreover, it gives the action of the dynamical semigroup $\tilde{\Phi}_t$ generated by (3.10) on any polynomial in the noncommutative variables p and q . Because $\tilde{\Phi}_t(AB) \neq \tilde{\Phi}_t(A)\tilde{\Phi}_t(B)$ it is not sufficient to know $\tilde{\Phi}_t(p)$ and $\tilde{\Phi}_t(q)$ as in the case of dynamical groups $\tilde{U}_t(AB) = \tilde{U}_t(A)\tilde{U}_t(B)$, where evidently it is sufficient to know the action of $\tilde{U}_t$ on p and q in order to know this action on all noncommutative polynomials. From theoretical point of view, the explicit action of $\tilde{\Phi}_t$ on Weyl operators $W(\xi, \eta)$ allows, as was shown by Lindblad in the Appendix of [16], directly to prove that the semigroup $\tilde{\Phi}_t$ is indeed a semigroup of completely positive mappings. This assertion cannot be considered as a consequence of the structural theorem of Lindblad because $\tilde{L}$ is unbounded.

We can write as Lindblad [16],

$$\tilde{\Phi}_t(W(\xi, \eta)) = W(\xi(t), \eta(t))e^{g(t)}$$

and then the master equation

$$\frac{d\tilde{\Phi}_t(W(\xi, \eta))}{dt} = \tilde{L}(\tilde{\Phi}_t(W(\xi, \eta)))$$

is equivalent with the following differential equations

$$\begin{aligned}\frac{d\xi(t)}{dt} &= -(\lambda + \mu)\xi(t) - \frac{1}{m}\eta(t), \\ \frac{d\eta(t)}{dt} &= m\omega^2\xi(t) - (\lambda - \mu)\eta(t), \\ \frac{dg(t)}{dt} &= -\frac{1}{\hbar^2}(D_{pp}\xi^2(t) + D_{qq}\eta^2(t) - 2D_{pq}\xi(t)\eta(t),\end{aligned}$$

where $\xi(0) = \xi$, $\eta(0) = \eta$ and $g(0) = 0$. The explicit determination of $\tilde{\Phi}_t(W(\xi, \eta))$ as a function of t reduces to the integration of these differential equations. We obtain [17]:

$$\xi(t) = \alpha(t)\xi(0) + \beta(t)\eta(0), \quad \eta(t) = \gamma(t)\xi(0) + \delta(t)\eta(0),$$

where

$$\begin{pmatrix} \alpha(t) & \beta(t) \\ \gamma(t) & \delta(t) \end{pmatrix} = e^{-\lambda t} \begin{pmatrix} \cosh \nu t - \frac{\mu}{\nu} \sinh \nu t & -\frac{1}{m\nu} \sinh \nu t \\ \frac{m\omega^2}{\nu} \sinh \nu t & \cosh \nu t + \frac{\mu}{\nu} \sinh \nu t \end{pmatrix}.$$

From

$$\tilde{\Phi}_t(W(\xi(0), \eta(0))) = W(\xi(t), \eta(t))e^{g(t)} = e^{(i/\hbar)(\eta(t)q - \xi(t)p) + g(t)}, \quad (3.29)$$

$$\frac{\partial W(\xi(0), \eta(0))}{\partial \xi(0)} \Big|_{\xi(0)=\eta(0)=0} = -\frac{i}{\hbar}p$$

and

$$\frac{\partial W(\xi(0), \eta(0))}{\partial \eta(0)} \Big|_{\xi(0)=\eta(0)=0} = \frac{i}{\hbar}q,$$

the action of $\tilde{\Phi}_t$ on q and p , respectively, is obtained:

$$\tilde{\Phi}_t(q) = \delta(t)q - \beta(t)p, \quad \tilde{\Phi}_t(p) = -\gamma(t)q + \alpha(t)p.$$

As an immediate consequence of these relations we have $[\tilde{\Phi}_t(q), \tilde{\Phi}_t(p)] = i\hbar I e^{-2i\lambda t}$. From (3.29) and

$$\begin{aligned}\frac{\partial^2 W(\xi(0), \eta(0))}{\partial \xi^2(0)} \Big|_{\xi(0)=\eta(0)=0} &= -\frac{1}{\hbar^2}p^2, \\ \frac{\partial^2 W(\xi(0), \eta(0))}{\partial \eta^2(0)} \Big|_{\xi(0)=\eta(0)=0} &= -\frac{1}{\hbar^2}q^2\end{aligned}$$

and

$$\frac{\partial^2 W(\xi(0), \eta(0))}{\partial \xi(0) \partial \eta(0)} \Big|_{\xi(0)=\eta(0)=0} = \frac{1}{2\hbar^2} (pq + qp),$$

it follows, respectively,

$$\tilde{\Phi}_t(q^2) = \tilde{\Phi}_t^2(q) + 2B(t),$$

$$\tilde{\Phi}_t(p^2) = \tilde{\Phi}_t^2(p) + 2A(t)$$

and

$$\tilde{\Phi}_t\left(\frac{pq + qp}{2}\right) = \tilde{\Phi}_t(p)\tilde{\Phi}_t(q) - 2C(t),$$

where

$$\begin{pmatrix} 2B(t)m\omega \\ 2A(t)/m\omega \\ -2C(t) \end{pmatrix} = T(e^{Kt} - I)K^{-1}TD.$$

Let us denote by $\sigma(t)$ the correlation matrix

$$\sigma(t) = \begin{pmatrix} m\omega\sigma_{qq}(t) & \sigma_{pq}(t) \\ \sigma_{pq}(t) & \sigma_{pp}(t)/m\omega \end{pmatrix}.$$

Now by definition, for any state $\rho \in \mathcal{D}(\mathcal{H})$:

$$\sigma_{qq}(t) = \text{Tr}(\rho\tilde{\Phi}_t(q^2)) - (\text{Tr}(\rho\tilde{\Phi}_t(q)))^2,$$

$$\sigma_{pp}(t) = \text{Tr}(\rho\tilde{\Phi}_t(p^2)) - (\text{Tr}(\rho\tilde{\Phi}_t(p)))^2$$

and

$$\sigma_{pq}(t) = \text{Tr}(\rho\tilde{\Phi}_t\left(\frac{pq + qp}{2}\right)) - \text{Tr}(\rho\tilde{\Phi}_t(p))\text{Tr}(\rho\tilde{\Phi}_t(q)).$$

Using the relations (3.20) and (3.21) the following interesting formulas are obtained

$$(\lambda^2 + \omega^2 - \mu^2) \det \sigma(\infty) = \frac{1}{4} \det \tilde{D} + \frac{1}{\lambda^2} \left(\frac{1}{2m} D_{pp} + \frac{m\omega^2}{2} D_{qq} + \mu D_{pq} \right)^2 \quad (3.30)$$

and

$$(\lambda^2 + \omega^2 - \mu^2) \det \sigma(\infty) = \frac{1}{4} \det \tilde{D} + \left(\frac{1}{2m} \sigma_{pp}(\infty) + \frac{m\omega^2}{2} \sigma_{qq}(\infty) + \mu \sigma_{pq}(\infty) \right)^2, \quad (3.31)$$

where

$$\tilde{D} = 2 \begin{pmatrix} m\omega D_{qq} & D_{pq} \\ D_{pq} & D_{pp}/m\omega \end{pmatrix}.$$

Comparison of (3.30) with (3.31) gives

$$\frac{1}{2m}\sigma_{pp}(\infty) + \frac{m\omega^2}{2}\sigma_{qq}(\infty) + \mu\sigma_{pq}(\infty) = \frac{1}{\lambda}\left(\frac{1}{2m}D_{pp} + \frac{m\omega^2}{2}D_{qq} + \mu D_{pq}\right). \quad (3.32)$$

But, the left-hand side of (3.32) is exactly the asymptotic mean value of the energy of the open harmonic oscillator. Hence (3.32) gives the value of $E(\infty)$ as a function of diffusion coefficients,

$$E(\infty) = \frac{1}{\lambda}\left(\frac{1}{2m}D_{pp} + \frac{m\omega^2}{2}D_{qq} + \mu D_{pq}\right).$$

Another expression for $E(\infty)$ which also follows from (3.30) and (3.31) is

$$E(\infty) = ((\lambda^2 + \omega^2 - \mu^2) \det \sigma(\infty) - \frac{1}{4} \det \tilde{D})^{1/2}.$$

The reality condition for $E(\infty)$ implies

$$\det \sigma(\infty) \geq \frac{1}{4(\lambda^2 + \omega^2 - \mu^2)} \det \tilde{D} \geq \frac{\lambda^2 \hbar^2}{4(\lambda^2 + \omega^2 - \mu^2)}$$

which is more restrictive for the overdamped case ($\omega < \mu$) than the generalized uncertainty inequality $\det \sigma(\infty) \geq \hbar^2/4$.

3.2 The method of the characteristic function

Instead of solving the master equation (3.7) directly, we first introduce the normally ordered quantum characteristic function $\chi(\Lambda, \Lambda^*, t)$ defined in terms of the density operator ρ by

$$\chi(\Lambda, \Lambda^*, t) = \text{Tr}[\rho(t) \exp(\Lambda a^+) \exp(-\Lambda^* a)], \quad (3.33)$$

where Λ is a complex variable and the trace is performed over the states of system. Substituting Eq.(3.33) into the master equation (3.7) and using the operator relations

$$a \exp(\Lambda a^+) = \exp(\Lambda a^+)(a + \Lambda), \quad a^+ \exp(-\Lambda^* a) = \exp(-\Lambda^* a)(a^+ + \Lambda^*),$$

$$\text{Tr}[\rho(t) \exp(\Lambda a^+) \exp(-\Lambda^* a)(a^+ + \Lambda^*)] = \partial_{\Lambda} \chi, \quad \text{Tr}[\rho(t) \exp(\Lambda a^+) \exp(-\Lambda^* a)a] = -\partial_{\Lambda^*} \chi$$

or applying the rules:

$$\rho \leftrightarrow \chi, \quad a\rho \leftrightarrow -\frac{\partial}{\partial \Lambda^*} \chi, \quad a^+ \rho \leftrightarrow \left(\frac{\partial}{\partial \Lambda} - \Lambda^*\right) \chi, \quad \rho a \leftrightarrow \left(-\frac{\partial}{\partial \Lambda^*} + \Lambda\right) \chi, \quad \rho a^+ \leftrightarrow \frac{\partial}{\partial \Lambda} \chi,$$

the following partial differential equation for χ is found [18]:

$$\begin{aligned} & \{\partial_t + [(\lambda - i\omega)\Lambda + \mu\Lambda^*]\partial_\Lambda + [(\lambda + i\omega)\Lambda^* + \mu\Lambda]\partial_{\Lambda^*}\}\chi(\Lambda, \Lambda^*, t) \\ & = \{L|\Lambda|^2 + C\Lambda^2 + C^*\Lambda^{*2}\}\chi(\Lambda, \Lambda^*, t), \end{aligned} \quad (3.34)$$

where

$$L = \lambda - D_2, \quad C = \frac{1}{2}(\mu + D_1^*).$$

We consider the state of the system initially to be a superposition of coherent states. Coherent states $|\alpha\rangle$ of the harmonic oscillator are minimum uncertainty states having the mean coordinate $\langle q \rangle$ and mean momentum $\langle p \rangle$ given by

$$\langle q \rangle = \langle \alpha | q | \alpha \rangle = \sqrt{\frac{2\hbar}{m\omega}} \text{Re}\alpha, \quad \langle p \rangle = \langle \alpha | p | \alpha \rangle = \sqrt{2\hbar m\omega} \text{Im}\alpha. \quad (3.35)$$

Consequently, we take as the initial density operator

$$\rho(0) = \int d\alpha d\beta N(\alpha, \beta) |\alpha\rangle\langle\beta|.$$

The quantum characteristic function corresponding to the operator $|\alpha\rangle\langle\beta|$ is given from Eq. (3.33) by

$$\chi = \langle \beta | \alpha \rangle \exp(\lambda\beta^* - \lambda^*\alpha). \quad (3.36)$$

We look for a solution of (3.34) having the exponential form

$$\chi(\Lambda, \Lambda^*, t) = \int d\alpha d\beta N(\alpha, \beta) \langle \beta | \alpha \rangle \exp[A(t)\Lambda + B(t)\Lambda^* + f(t)\Lambda^2 + f^*(t)\Lambda^{*2} + h(t)|\Lambda|^2]. \quad (3.37)$$

The form of the solution (3.37) is suggested from the fact that the left-hand side of Eq. (3.34) contains first-order derivatives with respect to the time and variables Λ and Λ^* and is symmetric with respect to complex conjugation. The functions $A(t)$, $B(t)$, $f(t)$ and $h(t)$ depend only on time. Corresponding to Eq. (3.36), these functions have to satisfy the initial conditions

$$A(0) = \beta^*, \quad B(0) = \alpha, \quad f(0) = 0, \quad h(0) = 0. \quad (3.38)$$

When we introduce the function (3.37) into Eq. (3.34) and equate the coefficients of equal powers of Λ and Λ^* , we get the following two systems of linear differential equations of first order with constant coefficients:

$$\frac{dA(t)}{dt} + (\lambda - i\omega)A(t) + \mu B(t) = 0$$

$$\frac{dB(t)}{dt} + \mu A(t) + (\lambda + i\omega)B(t) = 0 \quad (3.39a)$$

and

$$\begin{aligned} \frac{dR(t)}{dt} + 2\lambda R(t) + 2\omega I(t) + \mu h(t) &= \text{Re}C \\ \frac{dI(t)}{dt} + 2\lambda I(t) - 2\omega R(t) &= \text{Im}C \\ \frac{dh(t)}{dt} + 4\mu R(t) + 2\lambda h(t) &= L, \end{aligned} \quad (3.39b)$$

where $R(t) = \text{Re}f(t)$, $I(t) = \text{Im}f(t)$ with the initial conditions $R(0) = I(0) = h(0) = 0$. Subject to the initial conditions (3.38), the homogeneous system (3.39a) has the solution [18]:

$$A(t) = u(t)\beta^* - v(t)\alpha, B(t) = -u^*(t)\alpha + v(t)\beta^*, \quad (3.40)$$

where

$$\begin{aligned} u(t) &= \frac{1}{2}[\exp(-\mu_- t) + \exp(-\mu_+ t) + \frac{2i\omega}{\mu_- - \mu_+}(\exp(-\mu_+ t) - \exp(-\mu_- t))], \\ v(t) &= \frac{\mu}{\mu_- - \mu_+}(\exp(-\mu_- t) - \exp(-\mu_+ t)). \end{aligned} \quad (3.41)$$

The eigenvalues $\mu_{\pm}$ are given by

$$\mu_{\pm} = \lambda \pm \sqrt{\mu^2 - \omega^2}, \quad \gamma \equiv \sqrt{\mu^2 - \omega^2}. \quad (3.42)$$

The system (3.39b) has the eigenvalues $-2\lambda, -2(\lambda \pm \sqrt{\mu^2 - \omega^2}) = -2\mu_{\pm}$ and in order to integrate it we apply the same method as for the system (3.14) in the preceding Subsection.

We obtain:

$$\begin{aligned} f(t) &= \frac{P}{2\mu} \exp(-2\mu_+ t)(\gamma - i\omega) - \frac{N}{2\mu} \exp(-2\mu_- t)(\gamma + i\omega) - \frac{i\mu M}{2\omega} \exp(-2\lambda t) + f(\infty), \\ h(t) &= M \exp(-2\lambda t) + N \exp(-2\mu_- t) + P \exp(-2\mu_+ t) + h(\infty). \end{aligned} \quad (3.43)$$

Here $M, N, P, f(\infty)$ and $h(\infty)$ are constants given by [18]:

$$\begin{aligned} M &= \frac{\omega}{\lambda\gamma^2}(\mu\text{Im}C + \frac{\omega L}{2}), \quad N = \frac{\mu}{2\gamma^2(\lambda - \gamma)}(\gamma\text{Re}C - \omega\text{Im}C - \frac{\mu L}{2}), \\ P &= -\frac{\mu}{2\gamma^2(\lambda + \gamma)}(\gamma\text{Re}C + \omega\text{Im}C + \frac{\mu L}{2}) \end{aligned}$$

and the asymptotic values connected with the diffusion coefficients D_{qq} , D_{pp} and D_{pq} are:

$$\begin{aligned} R(\infty) &= \frac{2(\lambda \text{Re}C - \omega \text{Im}C) - L\mu}{4(\lambda^2 - \gamma^2)}, \\ I(\infty) &= \frac{2\omega\lambda \text{Re}C + 2(\lambda^2 - \mu^2)\text{Im}C - L\mu\omega}{4\lambda(\lambda^2 - \gamma^2)}, \\ h(\infty) &= \frac{L(\lambda^2 + \omega^2) - 2\mu(\lambda \text{Re}C - \omega \text{Im}C)}{2\lambda(\lambda^2 - \gamma^2)}. \end{aligned} \quad (3.44)$$

By knowing the characteristic function (3.37), (3.40)–(3.43) corresponding to the initial density operator which represents a superposition of coherent states, it is easy to obtain explicit formulae for the moments:

$$\langle a^{+m}(t)a^n(t) \rangle = \text{Tr}[a^{+m}(t)a^n(t)\rho(t)] = (-1)^n \frac{\partial^{n+m}}{\partial \Lambda^{*n} \partial \Lambda^m} \chi(\Lambda, \Lambda^*, t)|_{\Lambda=\Lambda^*=0}.$$

In the following, we take the density operator ρ in the coherent state representation

$$\rho(0) = \int P(\alpha) |\alpha\rangle \langle \alpha| d^2\alpha,$$

where $P(\alpha)$ is the diagonal or Glauber P distribution and $d^2\alpha = d\text{Re}\alpha d\text{Im}\alpha$. The integration covers the entire complex α plane. Then the characteristic function (3.37) becomes:

$$\chi(\Lambda, \Lambda^*, t) = \int d^2\alpha P(\alpha) \exp[(u\alpha^* - v\alpha)\Lambda + (-u^*\alpha + v\alpha^*)\Lambda^*] \exp[f\Lambda^2 + f^*\Lambda^{*2} + h|\Lambda|^2].$$

Let us assume that the damped oscillator is at $t = 0$ prepared in a pure coherent state, say $|\alpha_0\rangle$, corresponding to $P(\alpha) = \delta(\text{Re}\alpha - \text{Re}\alpha_0)\delta(\text{Im}\alpha - \text{Im}\alpha_0)$. One has

$$\chi^{(0)}(\Lambda, \Lambda^*, t) = \exp[(u\alpha_0^* - v\alpha_0)\Lambda + (-u^*\alpha_0 + v\alpha_0^*)\Lambda^*] \exp[f\Lambda^2 + f^*\Lambda^{*2} + h|\Lambda|^2].$$

The first moments are given by

$$\langle a^+(t) \rangle = \frac{\partial \chi^{(0)}(t)}{\partial \Lambda} \Big|_{\Lambda=\Lambda^*=0} = u\alpha_0^* - v\alpha_0, \quad \langle a(t) \rangle = -\frac{\partial \chi^{(0)}(t)}{\partial \Lambda^*} \Big|_{\Lambda=\Lambda^*=0} = u^*\alpha_0 - v\alpha_0^*.$$

Then, with the notations (3.9), using (3.35) and the transformations

$$q(t) = \sqrt{\frac{\hbar}{2m\omega}}(a^+(t) + a(t)), \quad p(t) = i\sqrt{\frac{\hbar m\omega}{2}}(a^+(t) - a(t))$$

for the displacement operator $q(t)$ and the momentum operator $p(t)$ of the oscillator, we obtain the following mean values:

$$\sigma_q(t) = \sqrt{\frac{\hbar}{2m\omega}}((u-v)\alpha_0^* + (u^* - v)\alpha_0), \quad \sigma_p(t) = i\sqrt{\frac{\hbar m\omega}{2}}((u+v)\alpha_0^* - (u^* + v)\alpha_0),$$

with u, v given by (3.41), (3.42). There are two cases:

a) the overdamped case: $\mu > \omega, \nu^2 = \mu^2 - \omega^2, \gamma \equiv \nu$; then

$$u(t) = \exp(-\lambda t)(\cosh \nu t + \frac{i\omega}{\nu} \sinh \nu t), \quad v(t) = -\frac{\mu}{\nu} \exp(-\lambda t) \sinh \nu t \quad (3.45)$$

and $\sigma_q(t), \sigma_p(t)$ take the previous form (3.15);

b) the underdamped case: $\mu < \omega, \Omega^2 = \omega^2 - \mu^2, \gamma \equiv i\Omega$; then

$$u(t) = \exp(-\lambda t)(\cosh \Omega t + \frac{i\omega}{\Omega} \sinh \Omega t), \quad v(t) = -\frac{\mu}{\Omega} \exp(-\lambda t) \sinh \Omega t \quad (3.46)$$

and $\sigma_q(t), \sigma_p(t)$ take the previous form (3.16).

For the variances one finds:

$$\begin{aligned} \langle a^2(t) \rangle &= \frac{\partial^2 \chi^{(0)}(t)}{\partial \Lambda^{*2}} \Big|_{\Lambda=\Lambda^*=0} = (u^* \alpha_0 - v \alpha_0^*)^2 + 2f^*, \\ \langle a^{+2}(t) \rangle &= \frac{\partial^2 \chi^{(0)}(t)}{\partial \Lambda^2} \Big|_{\Lambda=\Lambda^*=0} = (u \alpha_0^* - v \alpha_0)^2 + 2f, \\ \langle a^+(t) a(t) \rangle &= -\frac{\partial^2 \chi^{(0)}(t)}{\partial \Lambda \partial \Lambda^*} \Big|_{\Lambda=\Lambda^*=0} = (u \alpha_0^* - v \alpha_0)(u^* \alpha_0 - v \alpha_0^*) - h. \end{aligned} \quad (3.47)$$

Then the relations (3.9) will give us the explicit time dependence of the variances $\sigma_{qq}(t)$, $\sigma_{pp}(t)$, $\sigma_{pq}(t)$. The asymptotic values of these variances are given by the following expressions [18]:

$$\begin{aligned} \sigma_{qq}(\infty) &= \frac{\hbar}{m\omega} \left(f + f^* - h + \frac{1}{2} \right) \Big|_{t \rightarrow \infty}, \\ \sigma_{pp}(\infty) &= -\hbar m\omega \left(f + f^* + h - \frac{1}{2} \right) \Big|_{t \rightarrow \infty}, \\ \sigma_{pq}(\infty) &= i\hbar(f - f^*) \Big|_{t \rightarrow \infty}. \end{aligned}$$

With $f(\infty) = R(\infty) + iI(\infty)$ and by using the formulas (3.44) for $R(\infty), I(\infty), h(\infty)$, the asymptotic values of the variances take the same form (3.20) as in the preceding subsection, as expected.

With the relations (3.47), the expectation value of the energy operator can be calculated:

$$E(t) = \hbar\omega(\langle a^\dagger a \rangle + \frac{1}{2}) + \frac{i\hbar\mu}{2}(\langle a^{+2} \rangle - \langle a^2 \rangle).$$

3.3 Quasiprobability distributions for the damped harmonic oscillator

The methods of quasiprobabilities have provided technical tools of great power for the statistical description of microscopic systems formulated in terms of the density operator [50-53]. The first quasiprobability distribution was introduced by Wigner [54] in a quantum-mechanical context. In quantum optics the P representation introduced by Glauber [55,56] and Sudarshan [57] provided many practical applications of quasiprobabilities. The development of quantum-mechanical master equations was combined with the Glauber P representation to give a Fokker-Planck equation for the laser [58,59]. One useful way to study the consequences of the master equation for the one-dimensional damped harmonic oscillator is to transform it into equations for the c -number quasiprobability distributions associated with the density operator. The resulting differential equations of the Fokker-Planck type for the distribution functions can be solved by standard methods and observables directly calculated as correlations of these distribution functions. However, the Fokker-Planck equations do not always have positive-definite diffusion coefficients. In this case one can treat the problem with the generalized P distribution [51].

First we present a short summary of the theory of quasiprobability distributions [53,60]. For the master equation (3.7) of the harmonic oscillator, physical observables can be derived from the expectation values of polynomials of the annihilation and creation operators. The expectation values are determined by using the quantum density operator ρ . Usually one expands the density operator with the aid of coherent states, defined as eigenstates of the annihilation operator: $a|\alpha\rangle = \alpha|\alpha\rangle$. They are given in terms of the eigenstates of the harmonic oscillator as

$$|\alpha\rangle = \exp(-\frac{|\alpha|^2}{2}) \sum_{n=0}^{\infty} \frac{1}{\sqrt{n!}} \alpha^n |n\rangle,$$

with the normalization $|\langle\beta|\alpha\rangle|^2 = \exp(-|\alpha-\beta|^2)$. In order to solve the master equation (3.7) we represent the density operator ρ by a distribution function over a c -number phase space. The chosen distribution function, introduced in [60], is defined as follows:

$$\Phi(\alpha, s) = \frac{1}{\pi^2} \int \chi(\Lambda, s) \exp(\alpha\Lambda^* - \alpha^*\Lambda) d^2\Lambda, \quad (3.48)$$

with the characteristic function

$$\chi(\Lambda, s) = \text{Tr}[\rho D(\Lambda, s)],$$

where $D(\Lambda, s)$ is the displacement operator

$$D(\Lambda, s) = \exp(\Lambda a^+ - \Lambda^* a + \frac{1}{2}s|\Lambda|^2).$$

The interval of integration in Eq. (3.48) is the whole complex Λ plane. Because of

$$\delta^2(\alpha) = \frac{1}{\pi^2} \int \exp(\alpha \Lambda^* - \alpha^* \Lambda) d^2 \Lambda,$$

the distribution function $\Phi(\alpha, s)$ is the Fourier transform of the characteristic function. Since the density operator is normalized by $\text{Tr} \rho = 1$, one obtains the normalization of Φ :

$$\int \Phi(\alpha, s) d^2 \alpha = 1.$$

In this paper we restrict ourselves to distribution functions with the parameters $s = 1, 0$ and -1 . These distribution functions can be used to calculate expectation values of products of annihilation and creation operators. For that purpose we first expand the displacement operator in a power series of the operators a and a^+ :

$$D(\Lambda, s) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{\Lambda^m (-\Lambda^*)^n}{m!n!} \{a^{+m} a^n\}_s. \quad (3.49)$$

The parentheses with the index s indicate the special representations of the polynomials depending on s . For example for $n = m = 1$ we have the s -ordered operators:

$$\{a^+ a\}_{s=1} = a^+ a, \quad \{a^+ a\}_{s=0} = \frac{1}{2}(a^+ a + a a^+), \quad \{a^+ a\}_{s=-1} = a a^+. \quad (3.50)$$

Expectation values of the s -ordered operators can be calculated as follows:

$$\begin{aligned} \langle \{a^{+m} a^n\}_s \rangle &= \text{Tr}[\rho \{a^{+m} a^n\}_s] = \text{Tr}[\rho (\frac{\partial}{\partial \Lambda})^m (-\frac{\partial}{\partial \Lambda^*})^n D(\Lambda, s)|_{\Lambda=0}] \\ &= (\frac{\partial}{\partial \Lambda})^m (-\frac{\partial}{\partial \Lambda^*})^n \chi(\Lambda, s)|_{\Lambda=0} = \int (\alpha^*)^m \alpha^n \Phi(\alpha, s) d^2 \alpha. \end{aligned}$$

For the last step we apply the inverse relation to Eq. (3.48):

$$\chi(\Lambda, s) = \int \Phi(\alpha, s) \exp(\Lambda \alpha^* - \Lambda^* \alpha) d^2 \alpha.$$

In the following we discuss the distribution functions for $s = 1, 0$ and -1 in more detail. For $s = 1$ we obtain the Glauber P function [56,61], for $s = 0$ the Wigner function [54] and for $s = -1$ the Q function [60]. For $s = 1$ we have

$$D(\Lambda, 1) = \exp(\Lambda a^+) \exp(-\Lambda^* a).$$

Then the s ordering in Eq. (3.49) corresponds to normal ordering. Since the Glauber P function is the Fourier transform of the characteristic function

$$\chi_N(\Lambda) = \text{Tr}[\rho \exp(\Lambda a^+) \exp(-\Lambda^* a)] = \chi(\Lambda, 1),$$

it follows from Eq. (3.48) that the distribution $\Phi(\alpha, 1)$ is identical to the P function. This function is used for an expansion of the density operator in diagonal coherent state projection operators [55-57,62]:

$$\rho = \int P(\alpha) d^2\alpha |\alpha\rangle\langle\alpha|.$$

Calculating the expectation values of normally ordered operator products we obtain the relation

$$\langle a^{+m} a^n \rangle = \int (\alpha^*)^m \alpha^n \Phi(\alpha, 1) d^2\alpha = \int (\alpha^*)^m \alpha^n P(\alpha) d^2\alpha,$$

from which we again derive $P(\alpha) = \Phi(\alpha, 1)$. Despite the formal similarity with averaging procedure with a classical probability distribution, the function $P(\alpha)$ is not a true probability distribution. Because of the overcompleteness of the coherent states, the P function is not a unique, well-behaved positive function for all density operators.

Cahill [63] studied the P representation for density operators which represent pure states and found a narrow class of states for which the P representation exists. These states can be generated from a particular coherent state $|\alpha\rangle$ by the application of a finite number of creation operators. For example, for the ground state of the harmonic oscillator it is easy to show that $\chi_N(\Lambda) = 1$ for all Λ . In that case the P function becomes $P(\alpha) = \delta^2(\alpha)$. The delta function and its derivatives are examples of a class of generalized functions known as tempered distributions [61]. Also Cahill [64] introduced a representation of the density operator of the electromagnetic field that is suitable for all density operators and that reduces to the coherent state P representation when the latter exists. The representation has no singularities.

Sudarshan [57] offered a singular formula for the P representation in terms of an infinite series of derivatives of the delta function. From the mathematical point of view, such a series is usually not considered to be a distribution function [61,62].

For $s = -1$ we have

$$D(\Lambda, -1) = \exp(-\Lambda^* a) \exp(\Lambda a^+).$$

The s ordering corresponds to antinormal ordering. Because the Q function is the Fourier transform of the characteristic function

$$\chi_A(\Lambda) = \text{Tr}[\rho \exp(-\Lambda^* a) \exp(\Lambda a^+)] = \chi(\Lambda, -1),$$

it follows from Eq. (3.48) that the distribution $\Phi(\alpha, -1)$ is the Q function. It is given by the diagonal matrix elements of the density operator in terms of coherent states:

$$Q(\alpha) = \frac{1}{\pi} \langle \alpha | \rho | \alpha \rangle.$$

Though for all density operators the Q function is bounded, non-negative and infinitely differentiable, it has the disadvantage that not every positive Q function corresponds to a positive semidefinite Hermitian density operator. Evaluating moments is only simple in the Q representation for antinormally ordered operator products.

For $s = 0$, the distribution $\Phi(\Lambda, 0)$ becomes the Wigner function W . The latter function is defined as the Fourier transform of the characteristic function

$$\chi_S(\Lambda) = \text{Tr}[\rho \exp(\Lambda a^+ - \Lambda^* a)] = \chi(\Lambda, 0).$$

Because this characteristic function is identical to $\chi(\Lambda, 0)$, we conclude that $\Phi(\alpha, 0)$ is the Wigner function $W(\alpha)$. Therefore, the Wigner function can be used to calculate expectation values of symmetrically ordered operators:

$$\langle \{a^{+m} a^n\}_{s=0} \rangle = \int (\alpha^*)^m \alpha^n W(\alpha) d^2 \alpha.$$

The symmetrically ordered operators are the arithmetic average of $(m+n)!/m!n!$ differently ordered products of m factors of a^+ and n factors of a . An example for $m = n = 1$ is given in Eq. (3.50).

The Wigner function is a nonsingular, uniformly continuous function of α for all density operators and may in general assume negative values. It is related to the density operator as follows:

$$W(\alpha) = \frac{1}{\pi^2} \int d^2\Lambda \text{Tr}[\exp(\Lambda(a^+ - \alpha^*) - \Lambda^*(a - \alpha))\rho].$$

Also it can be obtained from the P representation:

$$W(\alpha) = \frac{2}{\pi} \int P(\beta) \exp(-2|\alpha - \beta|^2) d^2\beta.$$

By using the standard transformations [65-67], the master equation (3.7) can be transformed into a differential equation for a corresponding c -number distribution. Here we apply these methods to derive Fokker-Planck equations for the before mentioned distributions: Glauber P , Q and Wigner W distributions. Using the relations

$$\begin{aligned} \frac{\partial D(\Lambda, s)}{\partial \Lambda} &= [(s-1)\frac{\Lambda^*}{2} + a^+]D(\Lambda, s) = D(\Lambda, s)[(s+1)\frac{\Lambda^*}{2} + a^+], \\ \frac{\partial D(\Lambda, s)}{\partial \Lambda^*} &= [(s+1)\frac{\Lambda}{2} - a]D(\Lambda, s) = D(\Lambda, s)[(s-1)\frac{\Lambda}{2} - a], \end{aligned}$$

one can derive the following rules for transforming the master equation (3.7) into Fokker-Planck equations in the Glauber $P(s=1)$, $Q(s=-1)$ and Wigner $W(s=0)$ representations:

$$\begin{aligned} a\rho &\leftrightarrow (\alpha - \frac{s-1}{2}\frac{\partial}{\partial \alpha^*})\Phi, & a^+\rho &\leftrightarrow (\alpha^* - \frac{s+1}{2}\frac{\partial}{\partial \alpha})\Phi, \\ \rho a &\leftrightarrow (\alpha - \frac{s+1}{2}\frac{\partial}{\partial \alpha^*})\Phi, & \rho a^+ &\leftrightarrow (\alpha^* - \frac{s-1}{2}\frac{\partial}{\partial \alpha})\Phi. \end{aligned}$$

Applying these operator correspondences (repeatedly, if necessary), we find the following Fokker-Planck equations for the distributions $\Phi(\alpha, s)$ [19]:

$$\frac{\partial \Phi(\alpha, s)}{\partial t} = -(\frac{\partial}{\partial \alpha}d_\alpha + \frac{\partial}{\partial \alpha^*}d_\alpha^*)\Phi(\alpha, s) + \frac{1}{2}(\frac{\partial^2}{\partial \alpha^2}D_{\alpha\alpha} + \frac{\partial^2}{\partial \alpha^{*2}}D_{\alpha\alpha}^* + 2\frac{\partial^2}{\partial \alpha \partial \alpha^*}D_{\alpha\alpha^*})\Phi(\alpha, s). \quad (3.51)$$

Here, $\Phi(\alpha, s)$ is $P(s=1)$, $Q(s=-1)$ and $W(s=0)$. While the drift coefficients are the same for the three distributions, the diffusion coefficients are different:

$$d_\alpha = -(\lambda + i\omega)\alpha + \mu\alpha^*, \quad D_{\alpha\alpha} = D_1 + s\mu, \quad D_{\alpha\alpha^*} = D_2 - s\lambda.$$

The Fokker-Planck equation (3.51) can also be written in terms of real coordinates x_1 and x_2 defined by

$$\alpha = x_1 + ix_2 \equiv \sqrt{\frac{m\omega}{2\hbar}} < q > + i \frac{1}{\sqrt{2\hbar m\omega}} < p >, \quad \alpha^* = x_1 - ix_2 \quad (3.52).$$

as follows:

$$\frac{\partial \Phi(x_1, x_2)}{\partial t} = -\left(\frac{\partial}{\partial x_1} d_1 + \frac{\partial}{\partial x_2} d_2\right) \Phi(x_1, x_2) + \frac{1}{2} \left(\frac{\partial^2}{\partial x_1^2} D_{11}^{(s)} + \frac{\partial^2}{\partial x_2^2} D_{22}^{(s)} + 2 \frac{\partial^2}{\partial x_1 \partial x_2} D_{12}^{(s)} \right) \Phi(x_1, x_2), \quad (3.53)$$

with the new drift and diffusion coefficients given by

$$d_1 = -(\lambda - \mu)x_1 + \omega x_2, \quad d_2 = -\omega x_1 - (\lambda + \mu)x_2,$$

$$D_{11}^{(s)} = \frac{1}{\hbar} m\omega D_{qq} - \frac{s}{2}(\lambda - \mu), \quad D_{22}^{(s)} = \frac{1}{\hbar} \frac{D_{pp}}{m\omega} - \frac{s}{2}(\lambda + \mu), \quad D_{12}^{(s)} = \frac{1}{\hbar} D_{pq}.$$

We note that the diffusion matrix

$$\begin{pmatrix} D_{11}^{(s)} & D_{12}^{(s)} \\ D_{12}^{(s)} & D_{22}^{(s)} \end{pmatrix}$$

for the P distribution ($s = 1$) needs not to be positive definite.

Since the drift coefficients are linear in the variables x_1 and x_2 ($i = 1, 2$):

$$d_i = -\sum_{j=1}^2 A_{ij} x_j, \quad A_{ij} = -\frac{\partial d_i}{\partial x_j},$$

with

$$A = \begin{pmatrix} \lambda - \mu & -\omega \\ \omega & \lambda + \mu \end{pmatrix} \quad (3.54)$$

and the diffusion coefficients are constant with respect to x_1 and x_2 , Eq. (3.53) describes an Ornstein-Uhlenbeck process [68,69].

The solution of the Fokker-Planck equation (3.53) can immediately be written down provided that the diffusion matrix D is positive definite. However, the diffusion matrix in the Glauber P representation is not, in general, positive definite. For example, if

$$D_{11}^P D_{22}^P - (D_{12}^P)^2 < 0,$$

the P distribution does not exist as a well-behaved function. In this situation, the so-called generalized P distributions can be taken which are well-behaved, normal ordering functions

[51]. The generalized P distributions are nondiagonal expansions of the density operator in terms of coherent states projection operators. The Q and W distributions always exist; they are Gaussian functions if they are initially of Gaussian type.

From Eq. (3.53) one can directly derive the equations of motion for the expectation values of the variables x_1 and x_2 ($i = 1, 2$):

$$\frac{d\langle x_i \rangle}{dt} = - \sum_{j=1}^2 A_{ij} \langle x_j \rangle. \quad (3.55)$$

By using Eqs. (3.5), (3.52) and (3.55) we obtain the equations of motion for the expectation values $\sigma_q(t), \sigma_p(t)$ of the coordinate and momentum of the harmonic oscillator which are identical with those derived in the preceding two subsections by using the Heisenberg representation and the method of characteristic function, respectively (see Eqs. (3.13)).

The variances of the variables x_1 and x_2 are defined by the expectation values

$$\sigma_{ij} = \langle x_i x_j \rangle - \langle x_i \rangle \langle x_j \rangle, \quad i, j = 1, 2.$$

They are connected with the variances and covariance of the coordinate q and momentum p by

$$\sigma_{qq} = (2\hbar/m\omega)\sigma_{11}, \quad \sigma_{pp} = 2\hbar m\omega\sigma_{22}, \quad \sigma_{pq} = \langle (pq + qp) \rangle - \langle p \rangle \langle q \rangle = 2\hbar\sigma_{12}$$

and can be calculated with the help of the variances of the quasiprobability distributions ($i, j = 1, 2$):

$$\sigma_{ij}^{(s)} = \int x_i x_j \Phi(x_1, x_2, s) dx_1 dx_2 - \int x_i \Phi(x_1, x_2, s) dx_1 dx_2 \int x_j \Phi(x_1, x_2, s) dx_1 dx_2.$$

The following relations exist between the various variances:

$$\sigma_{ii} = \sigma_{ii}^P + \frac{1}{4} = \sigma_{ii}^Q - \frac{1}{4} = \sigma_{ii}^W, \quad i = 1, 2, \quad \sigma_{12} = \sigma_{12}^P = \sigma_{12}^Q = \sigma_{12}^W.$$

The variances $\sigma_{ij}^{(s)}$ fulfill the following equations of motion:

$$\frac{d\sigma_{ij}^{(s)}}{dt} = - \sum_{l=1}^2 (A_{il} \sigma_{lj}^{(s)} + \sigma_{il}^{(s)} A_{lj}^T) + D_{ij}^{(s)}. \quad (3.56)$$

They can be written explicitly in the form:

$$\begin{aligned}\frac{d\sigma_{11}^{(s)}}{dt} &= -2A_{11}\sigma_{11}^{(s)} - 2A_{12}\sigma_{12}^{(s)} + D_{11}^{(s)}, \\ \frac{d\sigma_{22}^{(s)}}{dt} &= -2A_{21}\sigma_{12}^{(s)} - 2A_{22}\sigma_{22}^{(s)} + D_{22}^{(s)}, \\ \frac{d\sigma_{12}^{(s)}}{dt} &= -(A_{11} + A_{22})\sigma_{12}^{(s)} - A_{21}\sigma_{11}^{(s)} - A_{12}\sigma_{22}^{(s)} + D_{12}^{(s)},\end{aligned}$$

where the matrix elements A_{ij} are defined in Eq. (3.54). These relations are sufficient to prove that the equations of motion for the variances σ_{11} and σ_{22} and the covariance σ_{12} are the same for all representations as expected. The corresponding equations of motion of the variances and covariance of the coordinate and momentum coincide with those obtained in the preceding two subsections by using the Heisenberg representation and the method of characteristic function, respectively (see Eqs. (3.14)).

In order that the system approaches a steady state, the condition $\lambda > \nu$ must be met. Thus the steady state solutions are

$$\Phi(x_1, x_2, s) = \frac{1}{2\pi\sqrt{\det(\sigma(\infty))}} \exp\left[-\frac{1}{2} \sum_{i,j=1,2} (\sigma^{-1})_{ij}(\infty) x_i x_j\right], \quad (3.57)$$

where the stationary covariance matrix

$$\sigma^{(s)}(\infty) = \begin{pmatrix} \sigma_{11}^{(s)}(\infty) & \sigma_{12}^{(s)}(\infty) \\ \sigma_{12}^{(s)}(\infty) & \sigma_{22}^{(s)}(\infty) \end{pmatrix}$$

can be determined from the algebraic equation (see Eq. (3.56)):

$$\sum_{l=1}^2 (A_{il}\sigma_{lj}^{(s)}(\infty) + \sigma_{il}^{(s)}(\infty)A_{lj}^T) = D_{ij}^{(s)}.$$

With the matrix elements A_{ij} given by (3.54), we obtain

$$\begin{aligned}\sigma_{11}^{(s)}(\infty) &= \frac{(2\lambda(\lambda + \mu) + \omega^2)D_{11}^{(s)} + \omega^2 D_{22}^{(s)} + 2\omega(\lambda + \mu)D_{12}^{(s)}}{4\lambda(\lambda^2 + \omega^2 - \mu^2)}, \\ \sigma_{22}^{(s)}(\infty) &= \frac{\omega^2 D_{11}^{(s)} + (2\lambda(\lambda - \mu) + \omega^2)D_{22}^{(s)} - 2\omega(\lambda - \mu)D_{12}^{(s)}}{4\lambda(\lambda^2 + \omega^2 - \mu^2)}, \\ \sigma_{12}^{(s)}(\infty) &= \frac{-\omega(\lambda + \mu)D_{11}^{(s)} + \omega(\lambda - \mu)D_{22}^{(s)} + 2(\lambda^2 - \mu^2)D_{12}^{(s)}}{4\lambda(\lambda^2 + \omega^2 - \mu^2)}.\end{aligned}$$

The explicit matrix elements $\sigma_{ij}^{(s)}$ for the three representations P, Q and W can be obtained by inserting the corresponding diffusion coefficients. The distribution functions (3.57) can be used to calculate the expectation values of the coordinate and momentum and the variances by direct integration. The following relations are noticed [48]:

$$\sigma_{ij}^W(\infty) = \frac{1}{2}(\sigma_{ij}^P(\infty) + \sigma_{ij}^Q(\infty)), \quad i, j = 1, 2.$$

3.4 The master equation in the Weyl-Wigner-Moyal representation

The Weyl-Wigner-Moyal representation is a remarkable phase-space representation of the quantum mechanics. Roughly speaking, a phase-space representation of the quantum mechanics is a mapping from the Hilbert space operators to the functions on the classical phase-space which is such that if A is mapped onto $f_A(x, y)$ and ρ is mapped onto $f_\rho(x, y)$, then

$$\text{Tr}(\rho A) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_\rho(x, y) f_A(x, y) dx dy. \quad (3.58)$$

In reality, it is not exactly so, because the Weyl mapping is a mapping from the functions on the phase-space to the Hilbert space operators. This mapping W was defined by Weyl [70] in the following way:

$$W(f) = \frac{1}{(2\pi\hbar)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(i/\hbar)(x\eta - y\xi)} f(x, y) dx dy \right) W(\xi, \eta) d\xi d\eta.$$

From this it follows very formally that for any $\rho \in \mathcal{D}(\mathcal{H})$,

$$\text{Tr}(\rho W(f)) = \frac{1}{(2\pi\hbar)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \left(\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(i/\hbar)(x\eta - y\xi)} \text{Tr}(\rho W(\xi, \eta)) d\xi d\eta \right) dx dy \quad (3.59)$$

and that (3.59) can be put in standard form (3.58) if the following function on the phase space is associated to any $\rho \in \mathcal{D}(\mathcal{H})$,

$$f_\rho(x, y) = \frac{1}{(2\pi\hbar)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(i/\hbar)(x\eta - y\xi)} \text{Tr}(\rho W(\xi, \eta)) d\xi d\eta. \quad (3.60)$$

The mapping $\rho \rightarrow f_\rho$ defined by (3.60) is exactly the Wigner mapping [54] which is the dual of the Weyl mapping $f \rightarrow W(f)$ (hence it can be denoted by $\widetilde{W}$) and $f_\rho = \widetilde{W}(\rho)$ is the Wigner function corresponding to the quantum state $\rho \in \mathcal{D}(\mathcal{H})$. But the quantum nature of the expectation value is not lost because $f_\rho(x, y)$ is not a probability distribution on the phase space, taking positive and negative values.

In the following, the phase space representation of the master equation (3.3) is obtained [17] by using the Wigner mapping (3.60). Denoting

$$f(x, y, t) = f_{\rho(t)}(x, y) = f_{\Phi_t(\rho)}(x, y),$$

it follows from the definition (3.60) that

$$f(x, y, t) = \frac{1}{(2\pi\hbar)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(i/\hbar)(x\eta - y\xi)} \text{Tr}(\rho(t)W(\xi, \eta)) d\xi d\eta. \quad (3.61)$$

Then

$$\frac{\partial f(x, y, t)}{\partial t} = \frac{1}{(2\pi\hbar)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(i/\hbar)(x\eta - y\xi)} \text{Tr}(L(\rho(t))W(\xi, \eta)) d\xi d\eta$$

and by duality

$$\frac{\partial f(x, y, t)}{\partial t} = \frac{1}{(2\pi\hbar)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(i/\hbar)(x\eta - y\xi)} \text{Tr}(\rho(t)\tilde{L}(W(\xi, \eta))) d\xi d\eta. \quad (3.62)$$

But from (3.10) in the case $t = 0$, $A = W(\xi, \eta)$ and by using the well-known identities for the Fourier transformation, Eq. (3.62) is transformed into the following evolution equation for the Wigner function [17]:

$$\begin{aligned} \frac{\partial f(x, y, t)}{\partial t} = & -\frac{y}{m} \frac{\partial f(x, y, t)}{\partial x} + m\omega^2 x \frac{\partial f(x, y, t)}{\partial y} + (\lambda - \mu) \frac{\partial}{\partial x} (xf(x, y, t)) \\ & + (\lambda + \mu) \frac{\partial}{\partial y} (yf(x, y, t)) + D_{qq} \frac{\partial^2 f(x, y, t)}{\partial x^2} + D_{pp} \frac{\partial^2 f(x, y, t)}{\partial y^2} + 2D_{pq} \frac{\partial^2 f(x, y, t)}{\partial x \partial y}. \end{aligned} \quad (3.63)$$

This equation looks classical and in fact is exactly an equation of the Fokker-Planck type. But not every function $f(x, y, 0)$ on the phase-space is the Wigner transform of a density operator. Hence, the quantum mechanics appears now in the restrictions imposed by this last condition on the initial condition $f(x, y, 0)$ for Eq. (3.63). Unfortunately, these restrictions are not known explicitly.

Because the most frequently used choice for $f(x, y, 0)$ is a Gaussian function and because Eq. (3.63) preserves this Gaussian type, i.e., $f(x, y, t)$ is also a Gaussian function, the differences between the quantum mechanics and classical mechanics are completely lost in this representation of the master equation. This is an explanation for the frequently occurring ambiguities on this subject in the literature.

Now from (3.61) by duality it follows that

$$f(x, y, t) = \frac{1}{(2\pi\hbar)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(i/\hbar)(x\eta - y\xi)} \text{Tr}(\rho(0) \tilde{\Phi}_t(W(\xi, \eta))) d\xi d\eta$$

and from the results of Subsection (3.1) it follows that

$$f(x, y, t) = \frac{1}{(2\pi\hbar)^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(i/\hbar)(x\eta - y\xi)} \text{Tr}(\rho(0) W(\xi(t), \eta(t)) e^{g(t)}) d\xi d\eta,$$

where

$$\begin{aligned} \xi(t) &= \alpha(t)\xi + \beta(t)\eta, \quad \eta(t) = \gamma(t)\xi + \delta(t)\eta, \\ \xi(0) &= \xi, \quad \eta(0) = \eta, \quad g(t) = -\frac{1}{\hbar^2} (A(t)\xi^2 + B(t)\eta^2 + 2C(t)\xi\eta). \end{aligned}$$

If the initial state $\rho(0)$ is a pure state corresponding to a coherent wave function

$$\rho(0)\phi = (\psi_{\sigma_q(0), \sigma_p(0)}, \phi) \psi_{\sigma_q(0), \sigma_p(0)}$$

centered in $x = \sigma_q(0)$, $y = \sigma_p(0)$, i.e., if

$$\psi_{\sigma_q(0), \sigma_p(0)}(x) = (W(\sigma_q(0), \sigma_p(0))\psi_0)(x),$$

where

$$\psi_0(x) = (2\pi\sigma_{qq}(0))^{-1/4} \exp(-x^2/4\sigma_{qq}(0)),$$

the Wigner function can be analytically calculated because the integrand is an exponential function with the exponent having a quadratic form in ξ and η .

With the help of the Wigner function the coordinate and momentum probability distribution are defined, respectively, by

$$P(x, t) = \int_{-\infty}^{\infty} f(x, y, t) dy$$

and

$$P(y, t) = \int_{-\infty}^{\infty} f(x, y, t) dx.$$

It follows

$$P(x, t) = \frac{1}{\sqrt{2\pi\sigma_{qq}(t)}} \exp\left(-\frac{(x - \sigma_q(t))^2}{2\sigma_{qq}(t)}\right)$$

and

$$P(y, t) = \frac{1}{\sqrt{2\pi\sigma_{pp}(t)}} \exp\left(-\frac{(y - \sigma_p(t))^2}{2\sigma_{pp}(t)}\right),$$

where, with the notations defined in Subsection (3.1)

$$\sigma_q(t) = \delta(t)\sigma_q(0) - \beta(t)\sigma_p(0),$$

$$\sigma_p(t) = \alpha(t)\sigma_p(0) - \gamma(t)\sigma_q(0),$$

$$\sigma_{qq}(t) = \delta^2(t)\sigma_{qq}(0) + \beta^2(t)\sigma_{pp}(0) + 2B(t),$$

$$\sigma_{pp}(t) = \alpha^2(t)\sigma_{pp}(0) + \gamma^2(t)\sigma_{qq}(0) + 2A(t)$$

and $\sigma_{pp}(0) = \hbar^2/4\sigma_{qq}(0)$. With these notations the Wigner function takes the following form [17]:

$$f(x, y, t) = \frac{1}{2\pi\sqrt{\sigma_{pp}(t)\sigma_{qq}(t) - \sigma_{pq}^2(t)}} \exp\left\{-\frac{1}{2(\sigma_{pp}(t)\sigma_{qq}(t) - \sigma_{pq}^2(t))}\right. \\ \left.\times (\sigma_{pp}(t)(x - \sigma_q(t))^2 + \sigma_{qq}(t)(y - \sigma_p(t))^2 - 2\sigma_{pq}(t)(x - \sigma_q(t))(y - \sigma_p(t)))\right\},$$

where

$$\sigma_{pq}(t) = \gamma(t)\delta(t)\sigma_{qq}(0) + \alpha(t)\beta(t)\sigma_{pp}(0) + 2C(t).$$

Evidently

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y, t) dx dy = 1.$$

3.5 Density matrix of the damped harmonic oscillator

In this Subsection we explore the general results that follow from the master equation of the one-dimensional damped harmonic oscillator. Namely, we discuss the physically relevant solutions of the master equation, by using the method of the generating function. In particular, we provide extended solutions (including both diagonal and off-diagonal matrix elements) for different initial conditions [20].

The method used in the following follows closely the procedure of Jang [21]. Let us first rewrite the master equation (3.7) for the density matrix by means of the number representation. Specifically, we take the matrix elements of each term between different

number states denoted by $|n\rangle$, and using $a^+|n\rangle = \sqrt{n+1}|n+1\rangle$ and $a|n\rangle = \sqrt{n}|n-1\rangle$, we get

$$\begin{aligned}
\frac{d\rho_{mn}}{dt} = & -i\omega(m-n)\rho_{mn} + \lambda\rho_{mn} - (m+n+1)D_2\rho_{mn} \\
& -\sqrt{m+1}nD_1^*\rho_{m+1,n-1} - \sqrt{m(n+1)}D_1\rho_{m-1,n+1} \\
& +\frac{1}{2}\sqrt{(n+1)(n+2)}(D_1-\mu)\rho_{m,n+2} + \frac{1}{2}\sqrt{(m+1)(m+2)}(D_1^*-\mu)\rho_{m+2,n} \\
& +\frac{1}{2}\sqrt{m(m-1)}(D_1+\mu)\rho_{m-2,n} + \frac{1}{2}\sqrt{(n-1)n}(D_1^*+\mu)\rho_{m,n-2} \\
& +\sqrt{(m+1)(n+1)}(D_2+\lambda)\rho_{m+1,n+1} + \sqrt{mn}(D_2-\lambda)\rho_{m-1,n-1}.
\end{aligned} \tag{3.64}$$

Here, we have used the abbreviated notation $\rho_{mn} = \langle m|\rho(t)|n\rangle$. This master equation is complicated in form and in the indices involved. It comprises not only the density matrix in symmetrical forms, such as $\rho_{m\pm 1, n\pm 1}$, but also matrix elements in asymmetrical forms like $\rho_{m\pm 2, n}, \rho_{m, n\pm 2}$ and $\rho_{m\mp 1, n\pm 1}$. In order to solve Eq. (3.64) we use the method of a generating function which allows us to eliminate the variety of indices m and n implicated in the equation. When we define the double-fold generating function by

$$G(x, y, t) = \sum_{m, n} \frac{1}{\sqrt{m!n!}} x^m y^n \rho_{mn}(t), \tag{3.65}$$

the density matrix can be evaluated from the inverse relation of Eq. (3.65):

$$\rho_{mn}(t) = \frac{1}{\sqrt{m!n!}} \left(\frac{\partial}{\partial x}\right)^m \left(\frac{\partial}{\partial y}\right)^n G(x, y, t)|_{x=y=0}, \tag{3.66}$$

provided that the generating function is calculated beforehand. When we multiply both sides of Eq. (3.64) by $x^m y^n / \sqrt{m!n!}$ and sum over the result, we get the following linear partial differential equation of second order for $G(x, y, t)$:

$$\begin{aligned}
\frac{\partial}{\partial t} G(x, y, t) = & \{[-(i\omega + D_2)x - D_1^*y] \frac{\partial}{\partial x} + [-D_1x + (i\omega - D_2)y] \frac{\partial}{\partial y} + (D_2 + \lambda) \frac{\partial^2}{\partial x \partial y} \\
& + \frac{1}{2}[(D_1^* - \mu) \frac{\partial^2}{\partial x^2} + (D_1 - \mu) \frac{\partial^2}{\partial y^2}] + [\frac{1}{2}(D_1 + \mu)x^2 + \frac{1}{2}(D_1^* + \mu)y^2 + (D_2 - \lambda)(xy - 1)]\} G(x, y, t).
\end{aligned} \tag{3.67}$$

The form of the solution of the Fokker-Planck equations treated in [17] and [21] suggest us to look for the solution of Eq. (3.67) of the form

$$G(x, y, t) = \frac{1}{A} \exp\{xy - [B(x - C)^2 + D(y - E)^2 + F(x - C)(y - E)]/H\}, \tag{3.68}$$

where A, B, C, D, E, F and H are unknown functions of time which have to be determined. When we first substitute the expression (3.68) for $G(x, y, t)$ into Eq. (3.67) and equate the coefficients of equal powers of x, y and xy on both sides of the equation, we get the following differential equations for the functions A, B, D, F and H :

$$-\frac{1}{A} \frac{dA}{dt} = -(D_1^* - \mu) \frac{B}{H} - (D_1 - \mu) \frac{D}{H} - (D_2 + \lambda) \frac{F}{H} + 2\lambda, \quad (3.69)$$

$$\frac{d}{dt} \left(\frac{B}{H} \right) = 2(\lambda - i\omega) \frac{B}{H} - \mu \frac{F}{H} - \frac{1}{2} (D_1 - \mu) \frac{F^2}{H^2} - 2(D_2 + \lambda) \frac{FB}{H^2} - 2(D_1^* - \mu) \frac{B^2}{H^2}, \quad (3.70)$$

$$\frac{d}{dt} \left(\frac{D}{H} \right) = 2(\lambda + i\omega) \frac{D}{H} - \mu \frac{F}{H} - \frac{1}{2} (D_1^* - \mu) \frac{F^2}{H^2} - 2(D_2 + \lambda) \frac{DF}{H^2} - 2(D_1 - \mu) \frac{D^2}{H^2}, \quad (3.71)$$

$$\frac{d}{dt} \left(\frac{F}{H} \right) = 2\lambda \frac{F}{H} - (D_2 + \lambda) \frac{F^2}{H^2} - 2\mu \left(\frac{B}{H} + \frac{D}{H} \right) - 4(D_2 + \lambda) \frac{DB}{H^2} - 2(D_1^* - \mu) \frac{BF}{H^2} - 2(D_1 - \mu) \frac{DF}{H^2}. \quad (3.72)$$

In addition to these equations, we get for the functions C and E

$$2B \frac{dC}{dt} + F \frac{dE}{dt} = (-2(\lambda - i\omega)B + \mu F)C + (2\mu B - (\lambda + i\omega)F)E, \quad (3.73)$$

$$2D \frac{dE}{dt} + F \frac{dC}{dt} = (-2(\lambda + i\omega)D + \mu F)E + (2\mu D - (\lambda - i\omega)F)C. \quad (3.74)$$

The equations (3.73) and (3.74) can be reformulated in order to eliminate the functions B, D and F , provided $BD - F^2/4 \neq 0$. We obtain

$$\frac{dC}{dt} = -(\lambda - i\omega)C + \mu E, \quad (3.75)$$

$$\frac{dE}{dt} = -(\lambda + i\omega)E + \mu C. \quad (3.76)$$

The functions A, B, D, F and H are connected by the auxiliary condition that $\text{Tr} \rho$ is independent of time. The trace of ρ can be evaluated by summing the diagonal matrix elements ρ_{nn} given in Eq.(3.66) or directly by using the integral expression

$$\text{Tr} \rho = \sum_{n=0}^{\infty} \rho_{nn} = \frac{1}{(2\pi)^2} \int \exp(-k_1 k_2) \exp(ik_1 x + ik_2 y) G(x, y, t) dk_1 dk_2 dx dy.$$

We obtain with the generating function (3.68)

$$\text{Tr} \rho = \left(\frac{4A^2}{H^2} \left(\frac{F^2}{4} - BD \right) \right)^{-1/2}. \quad (3.77)$$

This quantity is time-independent which can be verified by constructing an equation satisfied by the quantity $(F^2/4 - BD)/H^2$. Combining Eqs. (3.70)–(3.72) we get

$$\frac{d}{dt}\left(\frac{F^2/4 - BD}{H^2}\right) = 2[2\lambda - (D_1^* - \mu)\frac{B}{H} - (D_1 - \mu)\frac{D}{H} - (D_2 + \lambda)\frac{F}{H}]\left(\frac{F^2/4 - BD}{H^2}\right).$$

We see immediately that the first factor on the right-hand side of this equation is identical with the right-hand side of Eq. (3.69). Accordingly, we find

$$\frac{d}{dt}\left(\left(\frac{F^2}{4} - BD\right)\frac{A^2}{H^2}\right) = 0.$$

Since the scaling function H is arbitrary, we simplify the following equations by the choice

$$\frac{F^2}{4} - BD = -H. \quad (3.78)$$

Setting $\text{Tr}\rho = 1$, we obtain from Eqs. (3.77) and (3.78) the normalization constant $A^2 = -H/4$. As a consequence, we can simplify Eqs. (3.70)–(3.72) by eliminating the function H from these equations. The resulting three equations are

$$\begin{aligned} \frac{dB}{dt} &= -2(\lambda + i\omega)B - \mu F + 2(D_1 - \mu), \\ \frac{dD}{dt} &= -2(\lambda - i\omega)D - \mu F + 2(D_1^* - \mu), \\ \frac{dF}{dt} &= -2\mu(B + D) - 2\lambda F - 4(D_2 + \lambda). \end{aligned} \quad (3.79)$$

These equations imply that the function D is complex conjugate to B , provided that the function F is real.

In order to integrate the equations for the time-dependent functions B, C, D, E and F we start with Eqs. (3.75) and (3.76). These equations imply that the function E is complex conjugate to the function C . By solving the coupled equations we find:

$$C(t) = E^*(t) = u(t)C(0) - v(t)C^*(0), \quad (3.80)$$

where $u(t)$ and $v(t)$ are given by (3.45) and (3.46) for the two considered cases: overdamped and underdamped, respectively. For integrating the system (3.79) we proceed in the same way as for integrating the system (3.14). With the assumption that F is real and

$$D(t) = B^*(t) = R(t) + iI(t),$$

we obtain explicitly:

$$\begin{aligned}
R(t) &= \frac{1}{2}(e^{-2\mu+t} + e^{-2\mu-t})\tilde{R} + \frac{1}{2\gamma}(e^{-2\mu+t} - e^{-2\mu-t})(\omega\tilde{I} + \frac{\mu}{2}\tilde{F}) + R(\infty), \\
I(t) &= e^{-2\lambda t}\frac{\mu}{\gamma^2}(\mu\tilde{I} + \frac{\omega}{2}\tilde{F}) \\
&\quad - \frac{\omega}{2\gamma^2}(e^{-2\mu+t} + e^{-2\mu-t})(\omega\tilde{I} + \frac{\mu}{2}\tilde{F}) - \frac{\omega}{2\gamma}(e^{-2\mu+t} - e^{-2\mu-t})\tilde{R} + I(\infty), \\
F(t) &= -e^{-2\lambda t}\frac{\omega}{\gamma^2}(2\mu\tilde{I} + \omega\tilde{F}) \\
&\quad + \frac{\mu}{\gamma^2}(e^{-2\mu+t} + e^{-2\mu-t})(\omega\tilde{I} + \frac{\mu}{2}\tilde{F}) + \frac{\mu}{\gamma}(e^{-2\mu+t} - e^{-2\mu-t})\tilde{R} + F(\infty),
\end{aligned}$$

where we used the notations:

$$\mu_{\pm} = \lambda \pm \gamma, \quad \gamma \equiv \sqrt{\mu^2 - \omega^2}, \quad \tilde{R} = R(0) - R(\infty), \quad \tilde{I} = I(0) - I(\infty), \quad \tilde{F} = F(0) - F(\infty).$$

We can also obtain the connection between the asymptotic values of $B(t), D(t), F(t)$ and the coefficients D_1, D_2, μ and λ :

$$\begin{aligned}
R(\infty) = \text{Re}D(\infty) &= \frac{\lambda(\text{Re}D_1 - \mu) + \omega\text{Im}D_1 + \mu(D_2 + \lambda)}{\lambda^2 - \gamma^2}, \\
I(\infty) = \text{Im}D(\infty) &= \frac{\omega\lambda(\text{Re}D_1 - \mu) + (\mu^2 - \lambda^2)\text{Im}D_1 + \omega\mu(D_2 + \lambda)}{\lambda(\lambda^2 - \gamma^2)}, \\
F(\infty) &= -2\frac{\mu[\lambda(\text{Re}D_1 - \mu) + \omega\text{Im}D_1] + (\lambda^2 + \omega^2)(D_2 + \lambda)}{\lambda(\lambda^2 - \gamma^2)}.
\end{aligned}$$

When all explicit expressions for A, B, C, D, E, F and H are introduced into Eq. (3.68), we obtain an analytical form of the generating function $G(x, y, t)$ which allows us to evaluate the density matrix.

If the constants involved in the generating function satisfy the relations

$$C(0) = 0, \quad R(0) = R(\infty), \quad I(0) = I(\infty), \quad F(0) = F(\infty),$$

we obtain the stationary solution

$$C(t) = E(t) = 0, \quad R(t) = R(0), \quad I(t) = I(0), \quad F(t) = F(0)$$

and

$$D(t) = B^*(t) = R(0) + iI(0), \quad H(t) = -4A^2(t) = R^2(0) + I^2(0) - F^2(0)/4.$$

Then the stationary solution of Eq. (3.67) is

$$G(x, y, t) = \frac{1}{A} \exp\left\{\left(1 - \frac{F}{H}\right)xy - \frac{1}{H}(Bx^2 + B^*y^2)\right\}. \quad (3.81)$$

In addition, for a thermal bath [17] with

$$m\omega D_{qq} = \frac{D_{pp}}{m\omega}, \quad D_{pq} = 0, \quad \mu = 0,$$

the stationary generating function is simply given by

$$G(x, y) = \frac{2\lambda}{D_2 + \lambda} \exp\left(\frac{D_2 - \lambda}{D_2 + \lambda}xy\right).$$

The same generating function can be found for large times, if the asymptotic state is a Gibbs state with $\mu = 0$. In this case we obtain with Eq. (3.26) and $\mu = 0$

$$D_2 = \lambda \coth \frac{\hbar\omega}{2kT}$$

and

$$G(x, y) = (1 - \exp(-\frac{\hbar\omega}{kT})) \exp(\exp(-\frac{\hbar\omega}{kT})xy).$$

The density matrix can be calculated with Eq. (3.66) and yields the Bose-Einstein distribution

$$\rho_{mn}(\infty) = (1 - \exp(-\frac{\hbar\omega}{kT})) \exp(-\frac{n\hbar\omega}{kT}) \delta_{mn}.$$

A formula for the density matrix can be written down by applying the relation (3.66) to the generating function (3.68). We get

$$\begin{aligned} \rho_{mn}(t) &= \frac{\sqrt{m!n!}}{A} \exp[-(BC^2 + DE^2 + FCE)/H] \\ &\times \sum_{n_1, n_2, n_3=0} \frac{(1 - \frac{F}{H})^{n_3} (-\frac{B}{H})^{n_1} (-\frac{D}{H})^{n_2} (\frac{2BC}{H} + \frac{FE}{H})^{m-2n_1-n_3} (\frac{2DE}{H} + \frac{FC}{H})^{n-2n_2-n_3}}{n_1!n_2!n_3!(m-2n_1-n_3)!(n-2n_2-n_3)!}. \end{aligned} \quad (3.82)$$

In the case that the functions $C(t)$ and $E(t)$ vanish, the generating function has the form of Eq. (3.81). Then the elements of the density matrix with an odd sum $m + n$ are zero: $\rho_{mn} = 0$ for $m + n = 2k + 1$ with $k = 0, 1, 2, \dots$. The lowest non-vanishing elements are given with $\rho_{mn} = \rho_{nm}$ as

$$\rho_{00} = \frac{1}{A}, \quad \rho_{20} = -\frac{\sqrt{2}B}{AH}, \quad \rho_{11} = \frac{1}{A}\left(1 - \frac{F}{H}\right),$$

$$\rho_{22} = \frac{2BB^*}{AH^2} + \frac{1}{A}\left(-\frac{F}{H}\right)^2, \quad \rho_{31} = -\left(1 - \frac{F}{H}\right)\frac{\sqrt{6}B}{AH}, \quad \rho_{40} = \frac{\sqrt{6}B^2}{AH^2}.$$

It is also possible to choose the constants in such a way that the functions B and D vanish at time $t = 0$ and $F(0) = H(0)$. Then the density matrix (3.82) becomes at $t = 0$ ($E = C^*$):

$$\rho_{mn}(0) = \frac{1}{\sqrt{m!n!}}(C^*(0))^m(C(0))^n \exp(-|C(0)|^2). \quad (3.83)$$

This is the initial Glauber packet. The diagonal matrix elements of Eq. (3.83) represent a Poisson distribution used also in the study of multi-phonon excitations in nuclear physics. In the particular case when we assume

$$D_1 = \mu = 0, \quad D_2 = \lambda, \quad B(0) = D(0) = 0, \quad F(0) = H(0) = -4,$$

the differential equations (3.79) yield $B(t) = D(t) = 0$ and $F(t) = H(t) = -4$. Then the density matrix subject to the initial Glauber packet is (see also [21])

$$\rho_{mn}(t) = \frac{1}{\sqrt{m!n!}}(C^*(t))^m(C(t))^n \exp(-|C(t)|^2),$$

where $C(t)$ is given by Eq. (3.80).

4. Applications to nuclear equilibration processes

4.1 Charge equilibration in deep inelastic reactions

Over many years experimental data have been measured in the field of deep inelastic heavy ion collisions [71,72]. The characteristic feature of these collisions is the binary character of the system, i.e. the final fragments have nearly the same masses as the initial nuclei. Deep inelastic collisions are mainly described by the dynamics of selected collective degrees of freedom. These are the relative motion of the nuclei, mass and charge exchange, the neck degree of freedom and surface vibrations of the fragments [73]. Another important feature of these reactions is the dissipation of energy and angular momentum out of the collective degrees of freedom into the intrinsic or single-particle degrees of freedom.

In most of the approaches the collective degrees of freedom are chosen in an *a priori* way, guided by measured macroscopic variables which do not cover the complete space associated with all of them. Also, intrinsic degrees of freedom play a role. Thus, by taking into account the coupling between the actually treated and intrinsic degrees of freedom and other collective degrees of freedom, irreversible processes occur in the dynamics of the actually treated collective variables.

One of the most characteristic processes in deep inelastic reactions is the fast redistribution of neutrons and protons among the colliding nuclei in the early stage of the reaction called charge equilibration, neutron excess mode or isospin relaxation. The conditional variance of the charge distribution at fixed mass asymmetry has been largely used as a sensitive quantity to investigate the nature of the charge equilibration process.

There are two opposite approaches which have been developed in order to describe the charge equilibration mode, namely the quantum mechanical collective treatment and transport theories. The collective treatment stresses the coherent properties of the process and considers this mode as connected to the isovector dipole giant resonance of the composite system. An experimental indication for a quantum process is that the charge variance reaches a saturation value at large values of the total kinetic energy loss. The other opposite view relies on stochastic exchange processes between the colliding heavy nuclei.

In the following the charge equilibration mode is treated as a one-dimensional collective mode in the charge asymmetry degree of freedom. The damping of this collective mode is due to the coupling to the remaining collective and intrinsic degrees of freedom.

One method of introducing dissipation in a quantum mechanical description of deep inelastic collisions is to assume that the energy dissipation is similar to the loss of energy of a harmonic oscillator coupled with a large number of other harmonic oscillators. This mechanism can be simulated by a friction term of the Kostin type in the Schrödinger equation. According to this line, the role of collective motion and quantum fluctuations for charge and mass equilibration in deep inelastic reactions has been investigated [74,75].

Another method of treating dissipation in quantum mechanics, especially for the description of coupled collective modes, is the axiomatic method of Lindblad. In this method, as we said before, a simple dynamics for the subsystem of the explicitly treated collective degrees of freedom is chosen, namely a semigroup of transformations which introduces a preferred direction in time and, therefore, can describe an irreversible process.

In the present Subsection, Lindblad's theory for open quantum systems is applied to the problem of the damping of a single collective degree of freedom in deep inelastic collisions. We show the description of the charge equilibration mode by a damped harmonic oscillator in the framework of the Lindblad theory [17,18,22].

For a comparison of the theoretical results with experimental data we consider the charge equilibration in deep inelastic collisions of heavy ions. We can take over the solutions for the centroids and variances of a damped oscillator given by (3.15), (3.16), (3.19), (3.27), (3.28) for a comparison in order to illustrate our results. In [22] the charge equilibration of the systems $^{56}\text{Fe} + ^{209}\text{Bi}$, $^{56}\text{Fe} + ^{238}\text{U}$ and $^{98}\text{Mo} + ^{98}\text{Mo}$ was analysed within a collective description of the charge asymmetry degree of freedom. The charge asymmetry coordinate is defined as

$$\xi = \frac{Z_1 - Z_2}{Z_1 + Z_2}$$

where Z_1 and Z_2 are the charge numbers of the fragments. The Hamiltonian used for the description of this degree of freedom is a harmonic oscillator Hamiltonian of the form

$$H_0 = \frac{P_\xi^2}{2B} + \frac{1}{2}K(\xi - \xi_0)^2.$$

The stiffness parameter K and mass B were calculated in [22] on the basis of the liquid drop model for the potential energy and by use of a semiempirical formula for the frequency $\omega = (K/B)^{1/2}$. For the system $^{56}\text{Fe} + ^{209}\text{Bi}$ the following values were obtained [22]: $B = 1.6127 \times 10^{-40} \text{ MeV s}^2$ and $K = 1.6363 \times 10^4 \text{ MeV}$, corresponding to an oscillator energy $\hbar\omega = 6.63 \text{ MeV}$. Figs. 1–3 show the charge centroids and charge asymmetry variances as functions of time for the reaction of ^{56}Fe on ^{209}Bi at $E_{lab} = 465 \text{ MeV}$. The full dots are the experimental data of Breuer et al. [76]. Since the experimental data were obtained as functions of the total energy loss, Pop et al. [22] have related the data with the reaction times in order to compare them with calculations. As a model they used a parametrization of the deflection function in terms of relaxation times for the dissipation of the radial kinetic energy and relative angular momentum and for the development of deformations [77–79]. These times were fixed by reproducing the ridge of the double-differential cross-section $d^2\sigma/d\Omega dE$ plotted as a contour diagram in the total kinetic energy against scattering angle (Wilczynski plot). For short reaction times the kinetic energy loss is proportional to the reaction time ($\sim 30 \text{ MeV}/10^{-22}\text{s}$). Our comparison does not depend much on the specific assumptions of the model used to derive the reaction times from the measured total energy losses.

The "experimental" values of the variance in Fig. 3 are constructed from the experi-

mental values of $\sigma_{\xi\xi}$ by using the equation (see Eqs. (3.14)) [22]

$$\frac{d\sigma_{\xi\xi}}{dt} = -2(\Lambda - \mu)\sigma_{\xi\xi} + \frac{2}{B}\sigma_{\xi p_\xi} + 2D_{\xi\xi}$$

and applying the corresponding fitted values for the parameters Λ, μ and $D_{\xi\xi}$. The curves are calculated with the parameters ξ_0, Λ and μ fitted to the experimental data in Fig. 1 and in addition with $\omega_1 = -2m\omega D_{qq}/\hbar, \omega_2 = -2D_{pp}/m\hbar\omega$ and $\omega_3 = 2D_{qp}/\hbar$ fitted to the experimental variances in Figs. 2 and 3 (here $q = \xi, p = p_\xi, m = B$). The set of parameters ω_i is restricted by the condition (3.4) which now reads as

$$\omega_1\omega_2 - \omega_3^2 \geq \Lambda^2.$$

The full curves in Figs.1–3 represent the overdamped solution ($\omega = 1.0073 \times 10^{22} s^{-1} < \mu = 2.33 \times 10^{22} s^{-1}$) for the centroids and variances. This solution is the only one for which a good description of all the experimental data could be obtained. In the case of the underdamped solution no set of parameters could be found fitting simultaneously the data for the centroids and variances. Similar results were also obtained for the other considered systems $^{56}\text{Fe} + ^{238}\text{U}$ and $^{98}\text{Mo} + ^{98}\text{Mo}$ in [22].

The parameters of the theory are accepted as free quantities, fitted to the experimental data of the charge distributions. The overdamped solution with imposed fundamental constraints for the damped quantum oscillator succeeded to describe the experimental data within quite restricted ranges of the model parameters. Both the underdamped and the equilibrium solutions can not describe the data for the charge centroids and the isobaric variances.

The given treatment has the advantage that analytical solutions can be obtained for the damping of a harmonic oscillator. This is in contrast to the method when one introduces a friction term of the Kostin type into the Schrödinger equation and solves the resulting equation numerically. However, future investigations have to be carried out to interpret the fitted parameters of the theory from a microscopical point of view. For example, the parameters can be related to those obtained from a microscopical derivation of Fokker-Planck equations.

4.2 Charge and neutron equilibration and the open quantum system of two harmonic oscillators

In the present Subsection we extend the work of Gupta et al. [80] on the dynamics of the charge equilibration process in deep inelastic collisions and treat the damping of the proton and neutron asymmetry degrees of freedom with the method of Lindblad [23]. The charge and mass distribution in di-nuclear systems can be described with continuous coordinates of the charge and neutron asymmetries defined by

$$\eta_Z = \frac{Z_1 - Z_2}{Z_1 + Z_2}, \quad \eta_N = \frac{N_1 - N_2}{N_1 + N_2}.$$

Here, $Z_1(N_1)$ and $Z_2(N_2)$ are the total charge numbers (neutron numbers) on the left-hand and right-hand side of a plane through the neck of the di-nuclear system [73]. Without damping, the charge and neutron asymmetry degrees of freedom are described by a wavefunction $\psi(\eta_Z, \eta_N, t)$, which is the solution of the following time-dependent Schrödinger equation:

$$H(\eta_Z, \eta_N)\psi(\eta_Z, \eta_N, t) = i\hbar\partial\psi(\eta_Z, \eta_N, t)/\partial t,$$

where the Hamiltonian of the model is given as $(|k_{ZN}| \leq (k_Z k_N)^{1/2})$:

$$H(\eta_Z, \eta_N) = -\frac{\hbar}{2B_{ZZ}} \frac{\partial^2}{\partial \eta_Z^2} - \frac{\hbar^2}{2B_{NN}} \frac{\partial^2}{\partial \eta_N^2} + \frac{1}{2}k_Z \eta_Z^2 + \frac{1}{2}k_N \eta_N^2 - k_{ZN} \eta_Z \eta_N. \quad (4.1)$$

The Hamiltonian has the simple structure of two coupled oscillators in the coordinates η_Z and η_N in order to keep the time development of the wavefunction analytically solvable. The string constants of the potential can be calculated with the liquid drop model for the sticking configuration of the nuclei, i.e. for the relative distance $R = R_1 + R_2$, where R_1 and R_2 are the radii of the two colliding nuclei (for more details see [80]).

Hofmann et al. [39] have considered the coupling of the mass asymmetry coordinate to the charge asymmetry coordinate on the basis of a density operator formalism using a quantum master equation in a perturbative treatment. The coupling of the neutron and charge asymmetry coordinates has also been studied in the framework of a two-dimensional Fokker-Planck equation by Gross and Hartmann [81], Schröder et al. [82], Birkelund et al. [83] and Merchant and Nörenberg [84].

With the Hamiltonian (4.1) as an example, we study the damping of two coupled oscillators in the framework of the Lindblad theory. In order to have a formalism which is generally applicable, we give the following formulae in terms of the two general coordinates q_1 and q_2 instead of η_Z and η_N . We present the equation of motion of the open quantum

system of two oscillators in the Heisenberg picture. With this equation we derive the time dependence of the expectation values of the coordinates and momenta and their variances. The connection with the Wigner function and Weyl operator is also discussed. Finally, we demonstrate the time dependence of the various quantities for a simplified version of the model, where the decay constants can be calculated analytically.

If $\tilde{\Phi}_t$ is the dynamical semigroup describing the time evolution of the open quantum system in the Heisenberg picture, then the master equation is given for an operator A as follows [14,16,17]:

$$\frac{d\tilde{\Phi}_t(A)}{dt} = \tilde{L}(\tilde{\Phi}_t(A)) = \frac{i}{\hbar}[H, \tilde{\Phi}_t(A)] + \frac{1}{2\hbar} \sum_j (V_j^+ [\tilde{\Phi}_t(A), V_j] + [V_j^+, \tilde{\Phi}_t(A)] V_j). \quad (4.2)$$

The operators $H, V_j, V_j^+ (j = 1, 2, 3, 4)$ are taken to be functions of the basic observables of the two quantum oscillators. The coordinates q_k and the momenta p_k obey the usual commutation relations ($k, l = 1, 2$):

$$[q_k, p_l] = i\hbar\delta_{kl}, \quad [q_k, q_l] = [p_k, p_l] = 0.$$

In order to obtain an analytically solvable model, H is taken to be a polynomial of second degree in these basic observables and V_j, V_j^+ are taken to be polynomials of only first degree. Then in the linear space spanned by $q_k, p_k (k = 1, 2)$, there exist four linearly independent operators $V_{j=1,2,3,4}$:

$$V_j = \sum_{k=1}^2 a_{jk} p_k + \sum_{k=1}^2 b_{jk} q_k,$$

where $a_{jk}, b_{jk} \in \mathbb{C}$ with $j = 1, 2, 3, 4$. It yields

$$V_j^+ = \sum_{k=1}^2 a_{jk}^* p_k + \sum_{k=1}^2 b_{jk}^* q_k,$$

where a_{jk}^*, b_{jk}^* are the complex conjugates of a_{jk}, b_{jk} .

The Hamiltonian H is chosen in the form of two coupled oscillators

$$H = \sum_{k=1}^2 \left(\frac{1}{2m_k} p_k^2 + \frac{m_k \omega_k^2}{2} q_k^2 \right) + k_{12} p_1 p_2 + \frac{1}{2} \sum_{k_1, k_2=1}^2 \mu_{k_1, k_2} (p_{k_1} q_{k_2} + q_{k_2} p_{k_1}) + \nu_{12} q_1 q_2.$$

Here we use the following abbreviations ($k = 1, 2$) :

$$D_{q_k q_l} = D_{q_l q_k} = \frac{\hbar}{2} \text{Re}(\mathbf{a}_k^* \mathbf{a}_l), \quad D_{p_k p_l} = D_{p_l p_k} = \frac{\hbar}{2} \text{Re}(\mathbf{b}_k^* \mathbf{b}_l), \quad D_{q_k p_l} = D_{p_l q_k} = -\frac{\hbar}{2} \text{Re}(\mathbf{a}_k^* \mathbf{b}_l),$$

$$\alpha_{12} = -\alpha_{21} = -\text{Im}(\mathbf{a}_1^* \mathbf{a}_2), \quad \beta_{12} = -\beta_{21} = -\text{Im}(\mathbf{b}_1^* \mathbf{b}_2), \quad \lambda_{kl} = -\text{Im}(\mathbf{a}_k^* \mathbf{b}_l). \quad (4.3)$$

The scalar products are formed with the vectors $\mathbf{a}_k, \mathbf{b}_k$ and their complex conjugates $\mathbf{a}_k^*, \mathbf{b}_k^*$. The vectors have the components

$$\mathbf{a}_k = (a_{1k}, a_{2k}, a_{3k}, a_{4k}), \quad \mathbf{b}_k = (b_{1k}, b_{2k}, b_{3k}, b_{4k}).$$

Now, as a consequence of the definitions (4.3) of the phenomenological constants which appear in $\tilde{L}(A)$ and of the positivity of the matrix formed by the four vectors $\mathbf{a}_1, \mathbf{a}_2, \mathbf{b}_1, \mathbf{b}_2$, it follows that the principal minors of this matrix are positive or zero. This matrix is given by:

$$\frac{1}{2} \hbar \begin{pmatrix} \mathbf{a}_1^* \mathbf{a}_1 & \mathbf{a}_1^* \mathbf{a}_2 & \mathbf{a}_1^* \mathbf{b}_1 & \mathbf{a}_1^* \mathbf{b}_2 \\ \mathbf{a}_2^* \mathbf{a}_1 & \mathbf{a}_2^* \mathbf{a}_2 & \mathbf{a}_2^* \mathbf{b}_1 & \mathbf{a}_2^* \mathbf{b}_2 \\ \mathbf{b}_1^* \mathbf{a}_1 & \mathbf{b}_1^* \mathbf{a}_2 & \mathbf{b}_1^* \mathbf{b}_1 & \mathbf{b}_1^* \mathbf{b}_2 \\ \mathbf{b}_2^* \mathbf{a}_1 & \mathbf{b}_2^* \mathbf{a}_2 & \mathbf{b}_2^* \mathbf{b}_1 & \mathbf{b}_2^* \mathbf{b}_2 \end{pmatrix}$$

$$= \begin{pmatrix} D_{q_1 q_1} & D_{q_1 q_2} - i\hbar\alpha_{12}/2 & -D_{q_1 p_1} - i\hbar\lambda_{11}/2 & -D_{q_1 p_2} - i\hbar\lambda_{12}/2 \\ D_{q_2 q_1} - i\hbar\alpha_{21}/2 & D_{q_2 q_2} & -D_{q_2 p_1} - i\hbar\lambda_{21}/2 & -D_{q_2 p_2} - i\hbar\lambda_{22}/2 \\ -D_{p_1 q_1} + i\hbar\lambda_{11}/2 & -D_{p_1 q_2} + i\hbar\lambda_{21}/2 & D_{p_1 p_1} & D_{p_1 p_2} - i\hbar\beta_{12}/2 \\ -D_{p_2 q_1} + i\hbar\lambda_{12}/2 & -D_{p_2 q_2} + i\hbar\lambda_{22}/2 & D_{p_2 p_1} - i\hbar\beta_{21}/2 & D_{p_2 p_2} \end{pmatrix}. \quad (4.4)$$

For example, we can write one of the conditions obtained from the positivity of (4.4):

$$D_{q_1 q_1} D_{q_2 q_2} - D_{q_1 q_2}^2 \geq \frac{1}{4} \hbar^2 \alpha_{12}^2.$$

This inequality and the corresponding ones derived from (4.4) are constraints imposed on the phenomenological constants by the fact that $\tilde{\Phi}_t$ is a dynamical semigroup [14,16,17].

The time-dependent expectation values of self-adjoint operators A and B can be written with the density operator ρ , describing the initial state of the quantum system, as follows:

$$m_A(t) = \text{Tr}(\rho \tilde{\Phi}_t(A)), \quad \sigma_{AB}(t) = \frac{1}{2} \text{Tr}(\rho \tilde{\Phi}_t(AB + BA)).$$

In the following we denote the vector with the four components $m_{q_i}(t), m_{p_i}(t), i = 1, 2$, by $\mathbf{m}(t)$ and the following 4×4 matrix by $\sigma(t)$:

$$\sigma(t) = \begin{pmatrix} \sigma_{q_1 q_1} & \sigma_{q_1 q_2} & \sigma_{q_1 p_1} & \sigma_{q_1 p_2} \\ \sigma_{q_2 q_1} & \sigma_{q_2 q_2} & \sigma_{q_2 p_1} & \sigma_{q_2 p_2} \\ \sigma_{p_1 q_1} & \sigma_{p_1 q_2} & \sigma_{p_1 p_1} & \sigma_{p_1 p_2} \\ \sigma_{p_2 q_1} & \sigma_{p_2 q_2} & \sigma_{p_2 p_1} & \sigma_{p_2 p_2} \end{pmatrix}.$$

Then via direct calculation of $\tilde{L}(q_k)$ and $\tilde{L}(p_k)$ we obtain

$$\frac{d\mathbf{m}}{dt} = Y\mathbf{m}, \quad (4.5)$$

where

$$Y = \begin{pmatrix} -\lambda_{11} + \mu_{11} & -\lambda_{12} + \mu_{12} & 1/m_1 & -\alpha_{12} + k_{12} \\ -\lambda_{21} + \mu_{21} & -\lambda_{22} + \mu_{22} & \alpha_{12} + k_{12} & 1/m_2 \\ -m_1\omega_1^2 & \beta_{12} - \nu_{12} & -\lambda_{11} - \mu_{11} & -\lambda_{21} - \mu_{21} \\ -\beta_{12} - \nu_{12} & -m_2\omega_2^2 & -\lambda_{12} - \mu_{12} & -\lambda_{22} - \mu_{22} \end{pmatrix}. \quad (4.6)$$

From (4.5) it follows that

$$\mathbf{m}(t) = M(t)\mathbf{m}(0) = \exp(tY)\mathbf{m}(0), \quad (4.7)$$

where $\mathbf{m}(0)$ is given by the initial conditions. The matrix $M(t)$ has to fulfil the condition:

$$\lim_{t \rightarrow \infty} M(t) = 0. \quad (4.8)$$

In order that this limit exists, Y must have only eigenvalues with negative real parts.

By direct calculation of $\tilde{L}(q_k q_l)$, $\tilde{L}(p_k p_l)$ and $\tilde{L}(q_k p_l + p_l q_k)$, ($k, l = 1, 2$), we obtain

$$\frac{d\sigma}{dt} = Y\sigma + \sigma Y^T + 2D, \quad (4.9)$$

where D is the matrix of the diffusion coefficients

$$D = \begin{pmatrix} D_{q_1 q_1} & D_{q_1 q_2} & D_{q_1 p_1} & D_{q_1 p_2} \\ D_{q_2 q_1} & D_{q_2 q_2} & D_{q_2 p_1} & D_{q_2 p_2} \\ D_{p_1 q_1} & D_{p_1 q_2} & D_{p_1 p_1} & D_{p_1 p_2} \\ D_{p_2 q_1} & D_{p_2 q_2} & D_{p_2 p_1} & D_{p_2 p_2} \end{pmatrix}$$

and Y^T the transposed matrix of Y . The time-dependent solution of (4.9) can be written as

$$\sigma(t) = M(t)(\sigma(0) - \Sigma)M^T(t) + \Sigma, \quad (4.10)$$

where $M(t)$ is defined in (4.7). The matrix Σ is time independent and solves the static problem (4.9) ($d\sigma/dt = 0$):

$$Y\Sigma + \Sigma Y^T + 2D = 0. \quad (4.11)$$

Now we assume that the following limit exists for $t \rightarrow \infty$:

$$\sigma(\infty) = \lim_{t \rightarrow \infty} \sigma(t). \quad (4.12)$$

In this case it follows from (4.10) and (4.8):

$$\sigma(\infty) = \Sigma. \quad (4.13)$$

Inserting (4.13) into (4.10) we obtain the basic equations for our purposes:

$$\sigma(t) = M(t)(\sigma(0) - \sigma(\infty))M^T(t) + \sigma(\infty), \quad (4.14)$$

where

$$Y\sigma(\infty) + \sigma(\infty)Y^T = -2D. \quad (4.15)$$

Now we want to discuss the time dependence of the Wigner function. This function is defined as:

$$\begin{aligned} f(x_1, x_2, y_1, y_2, t) = & \frac{1}{(2\pi\hbar)^4} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp\left(-\frac{i}{\hbar}(x_1\eta_1 + x_2\eta_2 - y_1\xi_1 - y_2\xi_2)\right) \\ & \times \text{Tr}[\rho\tilde{\Phi}_t(W(\xi_1, \xi_2; \eta_1, \eta_2))] d\xi_1 d\xi_2 d\eta_1 d\eta_2, \end{aligned} \quad (4.16)$$

where the Weyl operator W is defined by $(\xi_1, \xi_2, \eta_1, \eta_2 \text{ real})$

$$W(\xi_1, \xi_2; \eta_1, \eta_2) = \exp[i\hbar^{-1}(\eta_1 q_1 + \eta_2 q_2 - \xi_1 p_1 - \xi_2 p_2)].$$

Using the method developed by Lindblad [14,16,17] for the one-dimensional case, we find the following relation for the time development of the Weyl operator:

$$\tilde{\Phi}_t(W(\xi_1, \xi_2; \eta_1, \eta_2)) = W(\xi_1(t), \xi_2(t); \eta_1(t), \eta_2(t)) \exp g(t). \quad (4.17)$$

The real functions $\xi(t) = (\xi_1(t), \xi_2(t); \eta_1(t), \eta_2(t))$ and $g(t)$ satisfy the equations of motion:

$$d\xi(t)/dt = JY^T J^{-1}\xi(t) \quad (4.18)$$

$$dg(t)/dt = -\hbar^{-2}\xi(t)JDJ^{-1}\xi(t), \quad (4.19)$$

where

$$J = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}.$$

Eqs. (4.18) and (4.19) are obtained by inserting the Weyl operator $W(\xi_1, \xi_2; \eta_1, \eta_2)$ into the equation of motion (4.2). The initial conditions for the coordinates $\xi_1(t), \xi_2(t), \eta_1(t)$ and $\eta_2(t)$ are determined by $\xi_1(0) = \xi_1, \xi_2(0) = \xi_2, \eta_1(0) = \eta_1$ and $\eta_2(0) = \eta_2$, respectively, and $g(t)$ by $g(0) = 0$. From (4.18) and (4.19) we find that $\xi(t)$ is a linear function in the coordinates ξ_1, ξ_2, η_1 and η_2 and $g(t)$ a quadratic function.

The Weyl operator can be used to calculate the time-dependent expectation values $\mathbf{m}(t)$ and $\sigma(t)$ (see (4.7) and (4.10)), since this operator is connected with the coordinates and momenta via the derivatives

$$\begin{aligned}\frac{\partial W}{\partial \xi_i} \Big|_{\xi=0} &= -\frac{i}{\hbar} p_i, \\ \frac{\partial W}{\partial \eta_i} \Big|_{\xi=0} &= \frac{i}{\hbar} q_i, \\ \frac{\partial^2 W}{\partial \xi_i \partial \xi_j} \Big|_{\xi=0} &= -\frac{1}{\hbar^2} p_i p_j, \\ \frac{\partial^2 W}{\partial \xi_i \partial \eta_j} \Big|_{\xi=0} &= \frac{1}{2\hbar^2} (p_i q_j + q_j p_i), \\ \frac{\partial^2 W}{\partial \eta_i \partial \eta_j} \Big|_{\xi=0} &= -\frac{1}{\hbar^2} q_i q_j.\end{aligned}\tag{4.20}$$

For example, one obtains by using (4.20)

$$\sigma_{p_i p_j}(t) = -\hbar^2 \text{Tr}(\rho \frac{\partial^2 \tilde{\Phi}_t(W)}{\partial \xi_i(0) \partial \xi_j(0)} \Big|_{\xi(t=0)=0}).$$

Equations of this type can be evaluated with the help of (4.17)–(4.19) and lead to the same results for $\mathbf{m}(t)$ and $\sigma(t)$ as given before. With the Weyl operator (4.17) we can calculate the time development of the Wigner function. For this purpose we use the Fourier transform of the Wigner function at $t = 0$:

$$\begin{aligned}& \text{Tr}\{\rho \exp[i\hbar^{-1}(\eta'_1 q_1 + \eta'_2 q_2 - \xi'_1 p_1 - \xi'_2 p_2)]\} \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp[i\hbar^{-1}(x_1 \eta'_1 + x_2 \eta'_2 - y_1 \xi'_1 - y_2 \xi'_2)] \\ & \quad \times f(x_1, x_2, y_1, y_2, t = 0) dx_1 dx_2 dy_1 dy_2.\end{aligned}$$

When this relation is inserted into (4.16) after the Weyl operator $\tilde{\Phi}_t(W)$ is expressed by (4.17), one can integrate over the coordinates ξ_1, ξ_2, η_1 and η_2 with the following result for the Wigner function:

$$f(x_1, x_2, y_1, y_2, t) = \frac{1}{[\det(4\pi Z)]^{1/2}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp[-\frac{1}{4}(\mathbf{x} - \mathbf{x}'M^T)Z^{-1}(\mathbf{x} - M\mathbf{x}')] \\ \times f(x'_1, x'_2, y'_1, y'_2, t=0) dx'_1 dx'_2 dy'_1 dy'_2, \quad (4.21)$$

where $\mathbf{x} = (x_1, x_2, y_1, y_2)$ and the matrix $Z(t)$ is given by

$$Z(t) = \int_0^t M(t')DM^T(t')dt'.$$

This definition can be applied in order to rewrite (4.14):

$$\sigma(t) = M(t)\sigma(0)M^T(t) + 2Z(t). \quad (4.22)$$

In the particular case when we set

$$f(x_1, x_2, y_1, y_2, t=0) = \frac{1}{[\det(2\pi\sigma(0))]^{1/2}} \exp[-\frac{1}{2}(\mathbf{x} - \mathbf{m}(0))\sigma(0)^{-1}(\mathbf{x} - \mathbf{m}(0))],$$

we obtain from (4.21):

$$f(x_1, x_2, y_1, y_2, t) = \frac{1}{[\det(2\pi\sigma(t))]^{1/2}} \exp[-\frac{1}{2}(\mathbf{x} - \mathbf{m}(t))\sigma(t)^{-1}(\mathbf{x} - \mathbf{m}(t))],$$

which is the well-known result for Wigner functions [45,69,85].

In order to illustrate the developed formalism we present an example of two oscillators, which are coupled by a potential of the form as used for the proton and neutron degrees of freedom in (4.1), i.e. $k_{12} = 0, \mu_{ij} = 0, \nu_{12} \neq 0$. In this case the matrix Y , governing the time development of the expectation values $\mathbf{m}(t)$ and $\sigma(t)$, becomes

$$Y = \begin{pmatrix} -\lambda_{11} & -\lambda_{12} & 1/m_1 & -\alpha_{12} \\ -\lambda_{21} & -\lambda_{22} & \alpha_{12} & 1/m_2 \\ -m_1\omega_1^2 & \beta_{12} - \nu_{12} & -\lambda_{11} & -\lambda_{21} \\ -\beta_{12} - \nu_{12} & -m_2\omega_2^2 & -\lambda_{12} & -\lambda_{22} \end{pmatrix}. \quad (4.23)$$

For the calculation of the matrix $M(t)$ we must diagonalize the matrix Y by solving the corresponding secular equation, i.e. $\det(Y - zI) = 0$, where z is the eigenvalue and I is the unit matrix. According to (4.23) one obtains an equation of fourth order for the

eigenvalues z , which can be simply solved only for special examples. In the particular case with $\alpha_{12} = 0, \beta_{12} = 0, \lambda_{12} = 0$ and $\lambda_{21} = 0$, the secular equation is obtained as

$$[(z + \lambda_{11})^2 + \omega_1^2][(z + \lambda_{22})^2 + \omega_2^2] = \nu_{12}^2/m_1m_2.$$

The roots of this equation have the general structure

$$z_1 = -\gamma_+ + i\omega_+, \quad z_2 = -\gamma_+ - i\omega_+, \quad z_3 = -\gamma_- + i\omega_-, \quad z_4 = -\gamma_- - i\omega_-.$$

The constants $\gamma_{\pm}$ and $\omega_{\pm}$ can be easily calculated for the case $\nu_{12} = 0$:

$$\gamma_+ = \lambda_{11}, \quad \gamma_- = \lambda_{22}, \quad \omega_+ = \omega_1, \quad \omega_- = \omega_2 \quad (4.24)$$

or for $\lambda_{11} = \lambda_{22} = \lambda, \omega_1 = \omega_2 = \omega (\omega^2 > \nu_{12}/(m_1m_2)^{1/2})$

$$\gamma_+ = \gamma_- = \lambda, \quad \omega_{\pm}^2 = \omega^2 \pm \nu_{12}/(m_1m_2)^{1/2}.$$

Only positive values of γ_+ and γ_- fulfil (4.8). Applying the eigenvalues z_i of Y we can write the time-dependent matrix $M(t)$ as follows:

$$M_{mn}(t) = \sum_i N_{mi} \exp(z_i t) N_{in}^{-1},$$

where the matrix N represents the eigenvectors of Y :

$$\sum_n Y_{mn} N_{ni} = z_i N_{mi}.$$

With the relations $M_{mn}(t=0) = \delta_{mn}$ and $dM_{mn}(t)/dt|_{t=0} = Y_{mn}$ and using (4.7), (4.14), we conclude that the expectation values of the coordinates and momenta decay with the exponential factors $\exp(-\gamma_+ t)$ and $\exp(-\gamma_- t)$ and the matrix elements σ_{mn} with the combined factors $\exp(-2\gamma_+ t)$, $\exp(-\gamma_- t)$ and $\exp[-(\gamma_+ + \gamma_-)t]$.

Since the matrix elements M_{mn} are in general lengthy expressions, we present here the matrix $M(t)$ only for the special and simple case that the oscillators are uncoupled. With the roots given in (4.24) we obtain

$$M(t) = \begin{pmatrix} e^{-\lambda_{11}t} \cos \omega_1 t & 0 & \frac{1}{m_1 \omega_1} e^{-\lambda_{11}t} \sin \omega_1 t & 0 \\ 0 & e^{-\lambda_{22}t} \cos \omega_2 t & 0 & \frac{1}{m_2 \omega_2} e^{-\lambda_{22}t} \sin \omega_2 t \\ -m_1 \omega_1 e^{-\lambda_{11}t} \sin \omega_1 t & 0 & e^{-\lambda_{11}t} \cos \omega_1 t & 0 \\ 0 & -m_2 \omega_2 e^{-\lambda_{22}t} \sin \omega_2 t & 0 & e^{-\lambda_{22}t} \cos \omega_2 t \end{pmatrix}. \quad (4.25)$$

This matrix can be used to evaluate $\sigma(t)$ defined by (4.14) or (4.22). For example, we find the following expression for $\sigma_{12} = \sigma_{q_1 q_2}$ with $M(t)$ of (4.25):

$$\begin{aligned}\sigma_{q_1 q_2}(t) = & \exp[-(\lambda_{11} + \lambda_{22})t][(\sigma_{q_1 q_2}(0) - \sigma_{q_1 q_2}(\infty)) \cos \omega_1 t \cos \omega_2 t \\ & + \frac{1}{m_1 \omega_1}(\sigma_{q_2 p_1}(0) - \sigma_{q_2 p_1}(\infty)) \sin \omega_1 t \cos \omega_2 t + \frac{1}{m_2 \omega_2}(\sigma_{q_1 p_2}(0) - \sigma_{q_1 p_2}(\infty)) \cos \omega_1 t \sin \omega_2 t \\ & + \frac{1}{m_1 m_2 \omega_1 \omega_2}(\sigma_{p_1 p_2}(0) - \sigma_{p_1 p_2}(\infty)) \sin \omega_1 t \sin \omega_2 t] + \sigma_{q_1 q_2}(\infty).\end{aligned}$$

Similar expressions are found for the other matrix elements of $\sigma(t)$. The matrix elements of $\sigma(\infty)$ depend on Y and D and must be evaluated with (4.15) or by the relation:

$$\sigma(\infty) = 2 \int_0^\infty M(t') D M^T(t') dt'.$$

As an example we present the value of $\sigma_{q_1 q_2}(\infty)$:

$$\begin{aligned}\sigma_{q_1 q_2}(\infty) = & 2\{[(\lambda_{11} + \lambda_{22})^2 + (\omega_1 + \omega_2)^2][(\lambda_{11} + \lambda_{22})^2 + (\omega_1 - \omega_2)^2]\}^{-1} \\ & \times \{(\lambda_{11} + \lambda_{22})[(\lambda_{11} + \lambda_{22})^2 + \omega_1^2 + \omega_2^2]D_{q_1 q_2} + [(\lambda_{11} + \lambda_{22})^2 + \omega_1^2 - \omega_2^2]D_{q_2 p_1}/m_1 \\ & + [(\lambda_{11} + \lambda_{22})^2 + \omega_2^2 - \omega_1^2]D_{q_1 p_2}/m_2 + 2(\lambda_{11} + \lambda_{22})D_{p_1 p_2}/m_1 m_2\}.\end{aligned}$$

Similar expressions are obtained for the other matrix elements of $\sigma(\infty)$. The diffusion coefficients $D_{q_1 q_2}$, $D_{q_1 p_2}$, $D_{q_2 p_1}$ and $D_{p_1 p_2}$ are in general zero for uncoupled oscillators interacting with an usual environment. This has the consequence that the expectation values $\sigma_{q_1 q_2}$, $\sigma_{q_1 p_2}$, $\sigma_{q_2 p_1}$ and $\sigma_{p_1 p_2}$ vanish for $t \rightarrow \infty$. It is a very interesting point that the general theory of Lindblad allows couplings via the environment between uncoupled oscillators with $k_{12} = 0$, $\mu_{ij} = 0$, $\nu_{12} = 0$. According to the definitions of the parameters in terms of the vectors $\mathbf{a}_k$ and $\mathbf{b}_k$, the diffusion coefficients above can be different from zero and can simulate an interaction between the "uncoupled" oscillators. In this case a structure of the environment is reflected in the motion of the oscillators.

5. Damping of angular momentum in open quantum systems

In heavy-ion collisions, especially in deep inelastic reactions, one has experimentally studied the loss of angular momentum in the relative motion of the nuclei [72]. Measurements of the γ -multiplicities yielded information on the spin of the excited nuclei, which means information on the loss of angular momentum in the relative motion of the nuclei.

If the energy loss in the relative motion is simultaneously measured one can determine a time scale of the loss of angular momentum, i.e. one can explore the damping process as a function of time. This example motivated us to study the damping of angular momentum. Another example for the damping of angular momentum is the spin-relaxation in the presence of a magnetic field which was theoretically studied in [86].

The damping of angular momentum can be described at different levels of approximations: one may use quantum-mechanical or classical methods and microscopical or phenomenological models. In this paper we describe the damping of angular momentum with the quantum-mechanical theory of Lindblad [24,25]. We choose the operators, which open the system, proportional to the components of the angular-momentum operator, to generators of the proper Lorentz group and proportional to a linear combination of momenta and coordinates.

Let us assume a finite set of Hermitian operators A_i depending on the basic variables of the physical problem and given in the Heisenberg picture. Then their time evolution can be determined within the theory of Lindblad by the following Markovian master equations:

$$\frac{dA_i(t)}{dt} = L(A_i(t)) = \sum_{k=1}^N M_{ik} A_k(t) + D_i, \quad i = 1, \dots, N.$$

The matrix elements M_{ik} and components D_i are time-independent numbers. They are calculated by the equations ($A_i = A_i(t=0)$):

$$L(A_i) = \frac{i}{\hbar} [H, A_i] + \frac{1}{2\hbar} \sum_j (V_j^+ [A_i, V_j] + [V_j^+, A_i] V_j) \quad (5.1).$$

Here, H is the Hamiltonian, V_j and V_j^+ are operators which are functions of the basic variables. V_j^+ is the Hermitian adjoint operator to V_j . The expectation values of the operators $A_i(t)$ are solutions of the differential equations ($i = 1, \dots, N$):

$$\frac{d\langle A_i(t) \rangle}{dt} = \sum_{k=1}^N M_{ik} \langle A_k(t) \rangle + D_i.$$

The following considerations are restricted to three spatial degrees of freedom described by the coordinates q_1, q_2, q_3 and linear momenta p_1, p_2, p_3 . We assume that the Hamiltonian conserves angular momentum:

$$[H, L_i] = 0, \quad i = 1, 2, 3.$$

Further we assume for simplicity that the opening of the quantum-mechanical system fulfils rotational symmetry. If we restrict this study to three operators $V_{j=1,2,3}$, these operators have to transform like the components of a vector under spatial rotations. They fulfil the commutators

$$[\sum_{j=1}^3 V_j^\dagger V_j, L_i] = 0, \quad i = 1, 2, 3.$$

The following three examples for V_j will be considered ($j = 1, 2, 3$) :

$$(1) \quad V_j = \alpha L_j,$$

$$(2) \quad V_j = \alpha N_j = \alpha(sp_j + q_j p_s),$$

$$(3) \quad V_j = ap_j + bq_j.$$

The operators N_j contain a fourth coordinate s and the corresponding momentum $p_s = -i\hbar\partial/\partial s$. This coordinate can be thought to describe an intrinsic degree of freedom not affected by spatial rotations. The operators L_j and N_j have commutation relations as the generators of the proper Lorentz group [87]. The free parameters α, a and b are complex numbers.

(1) For the case V_j proportional to L_j , we obtain from Eq. (5.1)

$$L(A) = \frac{i}{\hbar}[H, A] + \frac{|\alpha|^2}{2\hbar} \sum_{j=1}^3 [L_j, [A, L_j]].$$

If we set the operator A equal to the square of the angular momentum, $A = L^2$, we find conservation of the total angular momentum:

$$\frac{dL^2(t)}{dt} = 0, \quad \frac{d\langle L^2(t) \rangle}{dt} = 0 \quad (5.2)$$

Here $\langle L^2(t) \rangle$ denotes the expectation value of $L^2(t)$. The squares of the components of L^2 get equalized:

$$\frac{dL_i^2(t)}{dt} = -|\alpha|^2 \hbar (3L_i^2(t) - L^2), \quad i = 1, 2, 3. \quad (5.3)$$

The expectation value of L_i^2 is calculated as

$$\langle L_i^2(t) \rangle = \langle L_i^2 \rangle_{t=0} \exp(-3\hbar|\alpha|^2 t) + \frac{\langle L^2 \rangle}{3} (1 - \exp(-3\hbar|\alpha|^2 t)),$$

where $\langle L^2 \rangle$ is time independent as shown in Eq. (5.2). These results prove that the ansatz (1) can be used in applications when the square of the angular momentum is conserved, but the squares of the individual components are equalized with time. Examples are depolarization phenomena.

We note the fact that in the derivation of Eqs. (5.2) and (5.3) we only used the commutators between angular-momentum operators. Therefore, the results in this case are also true for systems with many degrees of freedom.

(2) For the case V_j proportional to N_j we obtain for $L(A)$ according to Eq. (5.1):

$$L(A) = \frac{i}{\hbar} [H, A] + \frac{|\alpha|^2}{2\hbar} \sum_{j=1}^3 [N_j, [A, N_j]].$$

If we choose $A = L^2$, we find the following time derivative of $L^2(t)$:

$$\frac{dL^2(t)}{dt} = 2|\alpha|^2 \hbar (L^2(t) + N^2(t)). \quad (5.4)$$

Here we used the commutation relations

$$[L_i, N_j] = i\hbar \sum_{l=1}^3 \varepsilon_{ijl} N_l, \quad [N_i, N_j] = -i\hbar \sum_{l=1}^3 \varepsilon_{ijl} L_l,$$

where ε_{ijl} is the Levi-Civita antisymmetric tensor. Since the right-hand side of Eq. (5.4) contains the square $N^2 = \sum_{j=1}^3 N_j^2$, we have also to calculate the time derivative of $N^2(t)$:

$$\frac{dN^2(t)}{dt} = \frac{i}{\hbar} [H, N^2(t)] + 2|\alpha|^2 \hbar (L^2(t) + N^2(t)). \quad (5.5)$$

In order to simplify the discussion of Eqs. (5.4) and (5.5) let us assume that the commutator of N^2 with H vanishes. Then the solution for the expectation values can be written as

$$\langle L^2(t) \rangle = \langle L^2 \rangle_{t=0} + \frac{1}{2} (\langle L^2 \rangle_{t=0} + \langle N^2 \rangle_{t=0}) (\exp(4|\alpha|^2 \hbar t) - 1), \quad (5.6)$$

$$\langle N^2(t) \rangle = \langle N^2 \rangle_{t=0} + \frac{1}{2} (\langle L^2 \rangle_{t=0} + \langle N^2 \rangle_{t=0}) (\exp(4|\alpha|^2 \hbar t) - 1). \quad (5.7)$$

This result shows that the expectation values of $L^2(t)$ and $N^2(t)$ increase exponentially with time. If the Hamiltonian H is a possitive function of L^2 and N^2 alone, also the energy would grow in time. In this case the master equation describes a physical system coupled

to a reservoir of infinite temperature which increases the energy and angular momentum exponentially. However, such a system is not applicable for a phenomenological description of nuclear processes. But it is thinkable that realistic applications in solid-state physics can be found.

Let us assume that the system has the smallest possible value of L^2 , namely $\langle L^2 \rangle_{t=0} = 0$. Then the increase of L^2 and N^2 is given according to Eqs. (5.6) and (5.7):

$$\langle L^2(t) \rangle = \frac{1}{2} \langle N^2 \rangle_{t=0} (\exp(4|\alpha|^2 \hbar t) - 1), \quad \langle N^2(t) \rangle = \frac{1}{2} \langle N^2 \rangle_{t=0} (\exp(4|\alpha|^2 \hbar t) + 1).$$

From these equations it is interesting to note that both $\langle L^2(t) \rangle$ and $\langle N^2(t) \rangle$ are directly proportional to $\langle N^2 \rangle_{t=0}$. Both expectation values are in general related by the equation

$$\langle L^2(t) \rangle - \langle L^2(t) \rangle = \langle N^2(t) \rangle - \langle N^2(t) \rangle_{t=0}.$$

(3) In the case $V_j = ap_j + bq_j$, we obtain

$$\begin{aligned} L(A) = & \frac{i}{\hbar} [H, A] + \frac{1}{\hbar^2} D_{pp} \sum_{j=1}^3 [q_j, [A, q_j]] + \frac{1}{\hbar^2} D_{qq} \sum_{j=1}^3 [p_j, [A, p_j]] \\ & - \frac{1}{\hbar^2} D_{pq} \sum_{j=1}^3 ([q_j, [A, p_j]] + [p_j, [A, q_j]]) + \frac{i\lambda}{2\hbar} \sum_{j=1}^3 ([q_j, [A, p_j]] + [A, p_j]q_j - p_j[A, q_j] - [A, q_j]p_j). \end{aligned} \quad (5.8)$$

In this equation we used the following abbreviations:

$$D_{qq} = \frac{\hbar}{2} a^* a, \quad D_{pp} = \frac{\hbar}{2} b^* b, \quad D_{pq} = -\frac{\hbar}{2} \text{Re}(a^* b), \quad \lambda = -\text{Im}(a^* b).$$

The quantities D_{qq} , D_{pp} and D_{pq} are denoted as the diffusion coefficients and λ as the friction constant. With the complex numbers a and b we can form the following determinant which is zero:

$$0 = \frac{\hbar}{2} \begin{vmatrix} a^* a & a^* b \\ b^* a & b^* b \end{vmatrix} = \begin{vmatrix} D_{qq} & -D_{pq} - i\hbar\lambda/2 \\ -D_{pq} + i\hbar\lambda/2 & D_{pp} \end{vmatrix}.$$

This determinant can be used to write down an equality serving as a constraint for the phenomenological parameters:

$$D_{qq}D_{pp} - D_{pq}^2 = \hbar^2 \lambda^2 / 4. \quad (5.9)$$

If we set $A = L^2$ in Eq. (5.8), we find

$$L(L^2) = 4(D_{pp}q^2 + D_{qq}p^2 - D_{pq}(\mathbf{qp} + \mathbf{pq})) - 4\lambda(L^2 + \frac{3}{2}\hbar^2), \quad (5.10)$$

with $q^2 = q_1^2 + q_2^2 + q_3^2$, $p^2 = p_1^2 + p_2^2 + p_3^2$ and $\mathbf{qp} = q_1p_1 + q_2p_2 + q_3p_3$. Since Eq. (5.10) contains other operators in addition to L^2 we have to calculate the corresponding equations of motion of these operators, too. We obtain

$$\frac{dq^2(t)}{dt} = \frac{i}{\hbar}[H, q^2(t)] - 2\lambda q^2(t) + 6D_{qq}, \quad (5.11)$$

$$\frac{dp^2(t)}{dt} = \frac{i}{\hbar}[H, p^2(t)] - 2\lambda p^2(t) + 6D_{pp}, \quad (5.12)$$

$$\frac{d(\mathbf{qp} + \mathbf{pq})_t}{dt} = \frac{i}{\hbar}[H, (\mathbf{qp} + \mathbf{pq})_t] - 2\lambda(\mathbf{qp} + \mathbf{pq})_t + 12D_{pq}. \quad (5.13)$$

Let us first assume that the Hamiltonian depends on L^2 only, e.g., $H = L^2/2\Theta$, where Θ is the moment of inertia. Then all commutators with H vanish in Eqs. (5.11)–(5.13). As function of time the expectation values of q^2 , p^2 and $\mathbf{qp} + \mathbf{pq}$ drop exponentially down ($\sim \exp(-2\lambda t)$). For large times they approach the final values

$$\langle q^2 \rangle_{t \rightarrow \infty} = 3D_{qq}/\lambda, \quad \langle p^2 \rangle_{t \rightarrow \infty} = 3D_{pp}/\lambda, \quad \frac{1}{2}\langle \mathbf{qp} + \mathbf{pq} \rangle_{t \rightarrow \infty} = 3D_{pq}/\lambda.$$

If we insert these values into the equation for $d\langle L^2(t) \rangle/dt$ using Eq. (5.10), we find for large times

$$\langle L^2 \rangle_{t \rightarrow \infty} = 0.$$

Solving the expectation values we obtain

$$\langle q^2(t) \rangle = A \exp(-2\lambda t) + 3D_{qq}/\lambda,$$

$$\langle p^2(t) \rangle = B \exp(-2\lambda t) + 3D_{pp}/\lambda,$$

$$\frac{1}{2}\langle (\mathbf{qp} + \mathbf{pq})_t \rangle = C \exp(-2\lambda t) + 3D_{pq}/\lambda,$$

$$\langle L^2(t) \rangle = D \exp(-4\lambda t) + \frac{2}{\lambda}(AD_{pp} + BD_{qq} - 2CD_{pq}) \exp(-2\lambda t).$$

The parameters A, B, C and D have to be chosen by initial conditions. A second example is the choice of the Hamiltonian of a spherical oscillator:

$$H = \frac{1}{2m}p^2 + \frac{m}{2}\omega^2 q^2.$$

If we calculate the expectation value of $L^2(t)$ for large times using the equations (5.10)–(5.13) we obtain

$$\langle L^2 \rangle_{t \rightarrow \infty} = \frac{6}{\lambda^2(\lambda^2 + \omega^2)} \left\{ \frac{1}{4} \left(\frac{D_{pp}}{m} - m\omega^2 D_{qq} \right)^2 + \omega^2 D_{pq}^2 \right\}. \quad (5.14)$$

This relation can be rewritten by using the expectation value of the energy at large times:

$$\langle H \rangle_{t \rightarrow \infty} = \frac{3}{2\lambda} \left(\frac{D_{pp}}{m} + m\omega^2 D_{qq} \right). \quad (5.15)$$

Inserting (5.15) into (5.14) and applying (5.9) we obtain

$$\langle L^2 \rangle_{t \rightarrow \infty} = \frac{2}{3(\lambda^2 + \omega^2)} (\langle H \rangle_{t \rightarrow \infty}^2 - (\frac{3}{2}\hbar\omega)^2).$$

If the diffusion coefficients are chosen in accordance with Eq. (5.9),

$$D_{pp} = \frac{\lambda\hbar}{2m\omega}, \quad D_{pp} = \frac{\lambda\hbar}{2}m\omega, \quad D_{pq} = 0,$$

we reach the lowest value of $\langle H \rangle_{t \rightarrow \infty} = (3/2)\hbar\omega$ and $\langle L^2 \rangle_{t \rightarrow \infty} = 0$.

In simple examples we have found operators V_j needed to open a system in which angular momentum is damped. The difficulty in choosing these operators is the fact that in general new and usually complex operators appear on the right-hand sides of the Lindblad equations after evaluating the commutators. For these new operators one has to determine their master equations. Only if the number of coupled master equations remains small, one can hope to find analytical solutions for the damping processes.

As shown in Eqs. (5.10)–(5.14) the damping depends on the parameter λ which results from a product of two free parameters. If only a single free parameter is contained in the operators V_j , as for example given in the case (2), we obtain no damping of the square of angular momentum in general but an exponential increase. Systems with an energy and angular momentum increasing in time are not suitable for application in nuclear physics.

A damping of angular momentum may be easier reached, if the rotational symmetry of the dissipative operators is given up. But the equations which have to be solved get more complex and cannot analytically be evaluated without approximations [88].

One can apply the damped three-dimensional harmonic oscillator presented here to heavy-ion collisions. For deep inelastic reactions of not too heavy nuclei one may approximate the internuclear potential near the barrier by a reverse parabola:

$$V(r) = V_B - \frac{m}{2}\kappa^2(r - r_B)^2. \quad (5.16)$$

Here, r is the internuclear distance and r_B the location of the barrier with height V_B . Equation (5.16) means a replacement of the frequency ω by $i\kappa$. A damping of angular momentum of the relative motion can be gained if the friction constant λ is larger than κ .

6. Quantum tunneling in open systems

Tunneling is a remarkable, essentially quantum phenomenon, consisting in the propagation of a particle through a classically impenetrable potential barrier. Standing at the basis of important applications in electronics, chemistry and nuclear physics, it has continuously been investigated from the beginning of the quantum mechanics [27],[89],[90], especially due to the difficulties raised by the effects of the dissipation [91-94], always present in practical cases.

In this Section, the master equation (3.1) is used for the study of dissipative phenomena on tunneling processes. We consider a fission potential barrier $U(q)$ [26], of the form represented in Fig. 4, where the first well corresponds to a compound nucleus and the second well to a fission channel. In order to generalize Gamow's formula of the tunneling rate, for the case when a dissipative environment is present we use the theory of perturbations. With the potential $U(q)$, the Hamiltonian is

$$H = \frac{p^2}{2m} + U(q) \quad (6.1)$$

and the master equation (3.1) becomes

$$\frac{d\rho}{dt} = -\frac{i}{\hbar}[H, \rho] + \Lambda(\rho), \quad (6.2)$$

where

$$\begin{aligned} \Lambda(\rho) = & -\frac{i\lambda}{2\hbar}([q, p\rho + \rho p] - [p, q\rho + \rho q]) \\ & -\frac{1}{\hbar^2}D_{qq}[p, [p, \rho]] - \frac{1}{\hbar^2}D_{pp}[q, [q, \rho]] + \frac{1}{\hbar^2}D_{pq}([q, [p, \rho]] + [p, [q, \rho]]). \end{aligned} \quad (6.3)$$

Using the commutation relation $[q, p] = i\hbar$, from Eqs. (6.2)–(6.3) we obtain the following evolution equations for the expectation values of the canonical coordinates q and p :

$$\begin{aligned} \frac{d\langle q \rangle}{dt} + \lambda \langle q \rangle &= \frac{\langle p \rangle}{m}, \\ \frac{d\langle p \rangle}{dt} + \lambda \langle p \rangle &= -\left\langle \frac{dU}{dq} \right\rangle \end{aligned} \quad (6.4)$$

and for their variances:

$$\begin{aligned}
\frac{d}{dt}\sigma_{qq} + 2\lambda\sigma_{qq} - \frac{2}{m}\sigma_{pq} &= 2D_{qq}, \\
\frac{d}{dt}\sigma_{pp} + 2\lambda\sigma_{pp} - 2\sigma_{pU} &= 2D_{pp}, \\
\frac{d}{dt}\sigma_{pq} + 2\lambda\sigma_{pq} - \frac{1}{m}\sigma_{pp} - \sigma_{qU} &= 2D_{pq},
\end{aligned} \tag{6.5}$$

where

$$\sigma_{qq} = \langle q^2 \rangle - \langle q \rangle^2, \quad \sigma_{pp} = \langle p^2 \rangle - \langle p \rangle^2, \quad \sigma_{pq} = \langle \frac{qp + pq}{2} \rangle - \langle q \rangle \langle p \rangle \tag{6.6}$$

and

$$\sigma_{pU} = -\frac{1}{2} \langle \frac{dU}{dq} p + p \frac{dU}{dq} \rangle + \langle \frac{dU}{dq} \rangle \langle p \rangle, \quad \sigma_{qU} = -\langle \frac{dU}{dq} q \rangle + \langle \frac{dU}{dq} \rangle \langle q \rangle. \tag{6.7}$$

In this way we see that the quantum master equation (6.2)–(6.3) leads to the Newtonian equations (6.4) with additional friction terms and to the similar equations (6.5) for the variances, with D_{qq}, D_{pp}, D_{pq} as zero-point values and $\sigma_{pq}, \sigma_{pU}, \sigma_{qU}$ as generalized forces.

Quantum tunneling is described as a transition from some "localized state" of the first well to some "localized state" of the second well. As we shall see later, this can be done by defining a Hamiltonian H_0 with approximate solutions in the first and in the second well, which disregard the tails corresponding to the other regions. The difference V between the two Hamiltonians defines a tunneling operator. Consequently, the Hamiltonian of the system takes the form:

$$H = H_0 + V, \tag{6.8}$$

where H_0 describes the localized states and V the transitions between them.

From Eqs. (6.4), with the expression $U(q) = -\omega_b^2 q^2/2, m = 1$ for the potential barrier, one obtains:

$$\frac{d \langle p \rangle}{d \langle q \rangle} = \frac{-\lambda \langle p \rangle + \omega_b^2 \langle q \rangle}{-\lambda \langle q \rangle + \langle p \rangle}, \tag{6.9}$$

which can be compared with the similar equation standing at the basis of the Dekker's theory of the quantum tunneling with dissipation [28], where the majority of the previous results are incorporated:

$$\frac{d \langle p \rangle}{d \langle q \rangle} = -2\lambda + \frac{\omega_b^2 \langle q \rangle}{\langle p \rangle}. \tag{6.10}$$

This equation has the solutions

$$\langle p \rangle_1 = (\sqrt{\omega_b^2 + \lambda^2} - \lambda) \langle q \rangle, \quad \langle p \rangle_2 = -(\sqrt{\omega_b^2 + \lambda^2} + \lambda) \langle q \rangle. \quad (6.11)$$

In this theory, the tunneling rate is considered of the form:

$$\Gamma(E) = \kappa \nu(E) P(E), \quad (6.12)$$

where $P(E)$ is the penetrability and the coefficient κ describing the effect of the dissipation is defined by the expression:

$$\kappa = \frac{\langle p(\lambda) \rangle}{\langle p(0) \rangle}. \quad (6.13)$$

With the first solution (6.11) describing the particle passing through the barrier, the expression (6.13) takes the explicit form:

$$\kappa = \sqrt{1 + \frac{\lambda^2}{\omega_b^2}} - \frac{\lambda}{\omega_b}, \quad (6.14)$$

which, in accordance to all previous results shows that due to the dissipation, the tunneling rate decreases. At the same time, with our expression (6.9) we obtain $\kappa = 1$. We conclude that the result (6.14) is a consequence of the particular way of introducing the dissipation in the equation (6.10), which is only empirically justified. When one considers the more accurate equations (6.4), resulting from the quantum theory of open systems and which can be easily understood from a physical point of view, the effect described by the expression (6.14) disappears. As we shall see in the following, the dissipative environment generates new transitions increasing the barrier penetrability.

By considering a basis of localized states, $|\Psi_0\rangle$ for the ground state of the first well (the compound nucleus) and $|\Psi_i\rangle$, $i = 1, 2, \dots$ for the second well (the reaction channel) the transition operator V takes the form:

$$V = \begin{pmatrix} 0 & V_{01} & V_{02} \dots & V_{0n} \\ V_{10} & 0 & 0 \dots & 0 \\ V_{20} & 0 & 0 \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ V_{n0} & 0 & 0 \dots & 0 \end{pmatrix}, \quad (6.15)$$

$$V_{i0} = V_{0i}, \quad i = 1, 2, \dots, n,$$

where the matrix elements have the approximate expressions [26]:

$$V_{0i} = V_{i0} = \frac{\hbar^2}{2m} [\Psi'_0(r_0)\Psi_i(r_0) - \Psi_0(r_0)\Psi'_i(r_0)]. \quad (6.16)$$

We consider that the operators V and $\Lambda(\rho)$ are small perturbations of the Hamiltonian H_0 defining the basis of the localized states. With the notations:

$$\rho'(t) = e^{\frac{i}{\hbar}H_0t}\rho(t)e^{-\frac{i}{\hbar}H_0t}, \quad V'(t) = e^{\frac{i}{\hbar}H_0t}V(t)e^{-\frac{i}{\hbar}H_0t}, \quad \Lambda'(\rho'(t);t) = e^{\frac{i}{\hbar}H_0t}\Lambda(\rho)e^{-\frac{i}{\hbar}H_0t} \quad (6.17)$$

in the interaction picture the quantum master equation (6.2) becomes:

$$\frac{d\rho'}{dt} = -\frac{i}{\hbar}[V'(t), \rho'(t)] + \Lambda'(\rho'(t);t). \quad (6.18)$$

Considering the zero-order density operator $\rho'^{(0)} = |0\rangle\langle 0|$, for the initial state the density operator in the second order approximation with respect to the operators V and Λ , can be written in the form:

$$\rho'(t) = \rho'^{(0)} + \rho'^{(1)}(t) + \rho'^{(2)}(t). \quad (6.19)$$

Introducing this expression in Eq. (6.18) and integrating, we obtain:

$$\rho'(t) = \rho'^{(0)} + \rho'_V(t) + \rho'_\Lambda(t) + \rho'_{VV}(t) + \rho'_{V\Lambda}(t) + \rho'_{LV}(t) + \rho'_{\Lambda\Lambda}(t), \quad (6.20)$$

where:

$$\begin{aligned} \rho'_V(t) &= -\frac{i}{\hbar} \int_0^t dt' [V'(t'), \rho'^{(0)}], \\ \rho'_\Lambda(t) &= \int_0^t dt' \Lambda'(\rho'^{(0)}; t'), \\ \rho'_{VV}(t) &= -\frac{1}{\hbar^2} \int_0^t dt' \int_0^{t'} dt'' [V'(t'), [V'(t''), \rho'^{(0)}]], \\ \rho'_{V\Lambda}(t) &= -\frac{i}{\hbar} \int_0^t dt' \int_0^{t'} dt'' [V'(t'), \Lambda'(\rho'^{(0)}; t'')], \\ \rho'_{\Lambda V}(t) &= -\frac{i}{\hbar} \int_0^t dt' \int_0^{t'} dt'' \Lambda'([V'(t''), \rho'^{(0)}]; t'), \\ \rho'_{\Lambda\Lambda}(t) &= \int_0^t dt' \int_0^{t'} dt'' \Lambda'(\Lambda'(\rho'^{(0)}; t''); t'). \end{aligned} \quad (6.21)$$

In principle we have to calculate the diagonal matrix elements $\rho_{ii}(t)$ in the representation of the localized states $|\psi_i(t)\rangle = \exp(-iH_0t/\hbar)|i\rangle, i = 1, 2, \dots$. In this case, $\rho_{ii}(t) = \langle \psi_i(t) | \rho | \psi_i(t) \rangle = \langle i | \rho'(t) | i \rangle$ and the transition rates have the expressions $\rho_{ii}(t)/t$. The functions $\rho_{ii}(t)/t$ describe the dependence of the transition probabilities on the frequency $\omega_i = (E_i - E_0)/\hbar$ due to the energy uncertainty corresponding to the time t . In this way we can see the influence of the openness on the tunneling spectrum and on the total tunneling rate $\Gamma = \sum_i \rho_{ii}(t)/t$. Using the expressions (6.3) and (6.15) of the openness and tunneling operators, neglecting the sums of the rapidly varying terms $\sum_j e^{i\omega_j t}$ and making the integrals (6.21), from the expression (6.20) we obtain:

$$\rho_{ii}(t) = \rho_{ii}^G(t) + \rho_{ii}^L(t) + \rho_{ii}^\lambda(t) + \rho_{ii}^D + \rho_{ii}^N + \rho_{ii}^A + \rho_{ii}^B(t) + \rho_{ii}^C(t) + \rho_{ii}^F(t) \quad (6.22)$$

with

$$\rho_{ii}^G(t) = \Omega_{i0}^2 \frac{\sin^2(\omega_i t/2)}{(\omega_i/2)^2}, \quad (6.23)$$

$$\rho_{ii}^L(t) = \left[\lambda C_{i0} \left(\lambda C_{0i} + 2D_{qq}u_{0i} - 2\frac{D_{pp}}{\hbar^2}w_{0i} \right) - 2\frac{D_{pq}}{\hbar}\Omega_{i0}(C_{0i} + 2v_{0i}) \right] \frac{\sin^2(\omega_i t/2)}{2(\omega_i/2)^2}, \quad (6.24)$$

$$\rho_{ii}^\lambda(t) = \lambda C_{i0}\Omega_{0i} \frac{\sin^2(\omega_i t/2)}{\omega_i/2} t, \quad (6.25)$$

$$\rho_{ii}^D(t) = -2(D_{qq}u_{00} + \frac{D_{pp}}{\hbar^2}w_{00})C_{i0}\Omega_{0i} \frac{t}{\omega_i} \left(\frac{\sin \omega_i t}{\omega_i t} - \cos \omega_i t \right), \quad (6.26)$$

$$\rho_{ii}^N(t) = -2\frac{D_{pq}}{\hbar}\lambda C_{i0}(C_{0i} + 2v_{0i}) \frac{t}{\omega_i} \left(1 - \frac{\sin \omega_i t}{\omega_i t} \right), \quad (6.27)$$

$$\rho_{ii}^A(t) = \left(-2\lambda q_{i0}s_{0i} + 2D_{qq}s_{0i}^2 + 2\frac{D_{pp}}{\hbar^2}q_{i0}^2 \right) t, \quad (6.28)$$

$$\rho_{ii}^B(t) = \left(-\lambda q_{i0}s_{0i} + D_{qq}s_{0i}^2 + \frac{D_{pp}}{\hbar^2}q_{i0}^2 \right) \lambda t^2, \quad (6.29)$$

$$\rho_{ii}^C = C_{i0}^2, \quad (6.30)$$

$$\begin{aligned} \rho_{ii}^F(t) = & C_{0i} \left\{ 2 \left[\Omega_{0i} - \frac{D_{pq}}{\hbar}(C_{0i} + 2v_{0i}) \right] \frac{\sin^2(\omega_i t/2)}{\omega_i/2} \right. \\ & \left. + \left(\lambda C_{0i} + 2D_{qq}u_{0i} - 2\frac{D_{pp}}{\hbar^2}w_{0i} \right) \frac{\sin \omega_i t}{\omega_i} \right\} \end{aligned} \quad (6.31)$$

depending on the following overlap integrals or matrix elements:

$$C_{0i} = C_{i0} = \int \Psi_0(q)\Psi_i(q)dq, \Omega_{0i} = \Omega_{i0} = \frac{V_{i0}}{\hbar},$$

$$\begin{aligned}
q_{i0} &= \int \Psi_0(q) q \Psi_i(q) dq, \quad s_{0i} = \int \Psi_0(q) d\Psi_i(q), \\
u_{0i} &= - \int \frac{d\Psi_0}{dq} \frac{d\Psi_i}{dq} dq, \quad u_{00} = \int \left(\frac{d\Psi_0}{dq} \right)^2 dq, \\
v_{0i} &= \int \Psi_0(q) q d\Psi_i(q), \quad w_{0i} = \int \Psi_0(q) q^2 \Psi_i(q) dq, \quad w_{00} = \int \Psi_0(q) q^2 \Psi_0(q) dq.
\end{aligned} \tag{6.32}$$

First of all, we notice that although the matrix elements given by the expressions (6.22)–(6.32) depend on the very small arbitrary distance $\delta\omega$ between the energy levels in the fission channel, the sum:

$$\sum_i \rho_{ii}(t) \simeq \int \rho_{ii}(\omega_i; t) \frac{d\omega_i}{\delta\omega} \tag{6.33}$$

of the probabilities on every span of energy do not depend on it, because every term contains two times the wave functions $\Psi_i(q)$ which, according to the expression (6.24) is proportional to $\sqrt{\delta\omega}$.

Secondly, one notice that, using the relation (6.33), the first term (6.23) leads to the Fermi's golden rule (6.32):

$$\Gamma^G = \sum_i \frac{\rho_{ii}^G(t)}{t} = 2\pi \frac{\Omega_{i0}^2}{\delta\omega}, \tag{6.34}$$

where we have neglected the dependence on i of the matrix element Ω_{i0} , in the very narrow span of energy $\Delta\omega = 1/t$. This term is symmetric with the energy $\hbar\omega_i = E_i - E_0$ and has a very small width of the spectrum, representing the energy uncertainty for the time t .

The second term (6.24), having the same time dependence as (6.23) is merely a correction to the Gamow's term, due to the openness parameters.

The third term (6.25), being asymmetric with ω_i , describes a tunneling energy shift proportional with the coefficient λ . Because the matrix element Ω_{i0} is negative, this term increases the transition rates for energies E_i smaller than the ground energy of the nucleus E_0 ($\omega_i < 0$), while for $E_i > E_0$ the transition rates are decreased.

The fourth term (6.26), rapidly varying with ω_i , describes transition rate modifications due to the diffusion coefficients D_{pp} and D_{qq} which, on the other hand, do not change the energy expectation values, the new introduced positive terms being compensated by negative ones in the summation.

The fifth term (6.27), depending on D_{pq} , introduces an energy shift similar to that introduced by λ . Because we have previously shown that at thermal equilibrium $D_{pq} = 0$ [17], this term describes nonequilibrium processes.

The sixth term (6.28) leads to a transition rate $\rho_{ii}^A(t)/t$ also not depending on time, but with a very large spectrum, having a dependence on ω_i much weaker than that of the functions multiplying the matrix elements in the expressions (6.23)–(6.27).

The seventh term (6.29) leads to a transition rate $\rho_{ii}^B(t)/t$ also with the very large spectrum given by the matrix elements q_{i0}, s_{0i} , but proportional with time. From the fundamental constraints (3.4) we find that both terms (6.28) and (6.29) are positive.

The terms (6.30) and (6.31) lead to a time-independent desintegration probability due to the overlap of the initial state $|0\rangle$ with the states $|i\rangle$.

For a long interval of time, the evolution of the system can be described by the expressions (6.22)–(6.31). In this case, the validity condition of the perturbation theory $\rho_{ii}(t) \ll \rho_{00}(0)$ holds also for large values of time, when $\sum_i \rho_{ii}(t) \gg \rho_{00}(0)$, the number of energy levels i being very large. Making the summation of the expressions (6.22)–(6.31), by using the formula (6.33), one obtains an expression of the form

$$\sum_i \rho_{ii}(t) = \chi + \Gamma_0 t + \Gamma_1 t^2, \quad (6.35)$$

where:

$$\begin{aligned} \Gamma_0 t &= \sum_i [\rho_{ii}^G(t) + \rho_{ii}^L(t) + \rho_{ii}^\lambda(t) + \rho_{ii}^D(t) + \rho_{ii}^N(t) + \rho_{ii}^A(t)], \\ \Gamma_1 t^2 &= \sum_i \rho_{ii}^B(t), \chi = \sum_i [C_{i0}^2 + \rho_{ii}^F(t)]. \end{aligned} \quad (6.36)$$

If $N(t)$ is the number of nondesintegrated nuclei, we can consider the equality

$$\sum_i \rho_{ii}(t) = - \int_0^t \frac{dN(t')}{N(t')}, \quad (6.37)$$

which leads to the following desintegration law :

$$N(t) = N_0 e^{-\chi - \Gamma_0 t - \Gamma_1 t^2} = N(0) e^{-\Gamma_0 t - \Gamma_1 t^2}. \quad (6.38)$$

In this expression appear two additional parameters: Γ_1 describing the irreversibility of the desintegration process in a dissipative environment and χ describing the nonorthogonality of the initial state $|0\rangle$ of the nucleus with the fission channel states $|i\rangle$. Estimating the time $t_0 = \chi/\Gamma_0 \sim m\Delta q^2/\hbar \sim 10^{-23}s$, where Δq is the width of the barrier, we find that χ can be neglected.

In the tunneling spectrum described by the expressions (6.22)–(6.31) we distinguish the Gamow's spectral line:

$$f(\omega_i t) = \frac{\sin^2(\omega_i t/2)}{(\omega_i t/2)^2} \quad (6.39)$$

represented in Fig. 5 and additional lines due to the openness, described by other functions:

$$g^\lambda(\omega_i t) = -\frac{\sin^2(\omega_i t/2)}{\omega_i t/2} \quad (6.40)$$

$$g^D(\omega_i t) = \frac{1}{\omega_i t} \left(\frac{\sin \omega_i t}{\omega_i t} - \cos \omega_i t \right), \quad (6.41)$$

$$g^N(\omega_i t) = -\frac{1}{\omega_i t} \left(1 - \frac{\sin \omega_i t}{\omega_i t} \right) \quad (6.42)$$

represented in Figs. 6, 7 and respectively 8. In this case, the transition rate spectrum can be written in the form:

$$\Gamma(\omega_i) = \frac{\rho_{ii}(t)}{t} = W_{i0}^2 t F(\omega_i t) + \varphi(\omega_i) \left(1 + \frac{\lambda}{2} t \right), \quad (6.43)$$

where:

$$W_{i0}^2 = \Omega_{i0}^2 + \lambda C_{i0} \left(\lambda C_{0i} + 2D_{qq}u_{0i} - 2\frac{D_{pp}}{\hbar^2}w_{0i} \right) - 2\frac{D_{pq}}{\hbar}\Omega_{i0}(C_{0i} + 2v_{0i}) \simeq \Omega_{i0}^2, \quad (6.44)$$

$$\varphi(\omega_i) = -2\lambda q_{i0}s_{0i} + 2D_{qq}s_{0i}^2 + 2\frac{D_{pp}}{\hbar^2}q_{i0}^2 \geq 0, \quad (6.45)$$

$$F(\omega_i t) = f(\omega_i t) + \eta^\lambda g^\lambda(\omega_i t) + \eta^D g^D(\omega_i t) + \eta^N g^N(\omega_i t), \quad (6.46)$$

the quantities

$$\begin{aligned} \eta^\lambda &= -\lambda \frac{C_{0i}\Omega_{0i}}{W_{i0}^2} \simeq -\lambda \frac{C_{i0}}{\Omega_{i0}} \geq 0, \\ \eta^D &= -2 \left(D_{qq}u_{00} + \frac{D_{pp}}{\hbar^2}w_{00} \right) \frac{C_{i0}\Omega_{0i}}{W_{i0}^2} \simeq -2 \left(D_{qq}u_{00} + \frac{D_{pp}}{\hbar^2}w_{00} \right) \frac{C_{i0}}{\Omega_{i0}} \geq 0, \\ \eta^N &= 2\frac{D_{pq}}{\hbar}\lambda \frac{C_{i0}(C_{0i} + 2v_{0i})}{W_{i0}^2} \geq 0 \end{aligned} \quad (6.47)$$

being practically independent, positive parameters, describing the four processes of the interaction of the system with the environment: the friction, the diffusion of q , the diffusion of p and respectively the non-equilibrium. The function $\varphi(\omega_i)$ is also positive due to the fundamental constraints.

7. Open quantum systems and the atom-field interaction

We consider a linearly polarized single mode electromagnetic field $\vec{E}$, of the frequency ω and the wavevector $\vec{k}_1$ propagating through an absorbing medium of two level atoms with the transition frequency ω_0 , the electric dipole μ and the density N ; every atom n is described by the Pauli operators $\sigma_x^n, \sigma_y^n, \sigma_z^n$. Due to the interaction between the radiation and the atoms, higher order harmonics $\vec{E}_\nu$ of the frequency $\nu\omega$ and wavevector $\vec{k}_\nu$ are generated. For the electric dipole interaction, the Hamiltonian of the system is of the form

$$H = -\frac{1}{2}\hbar\omega_0 \sum_n \sigma_z^n + \frac{1}{2} \sum_\nu (p_\nu^2 + \nu^2\omega^2 q_\nu^2) + \bar{\mu} \sum_{n\nu} \sigma_x^n (\nu\omega q_\nu \sin \vec{k}_\nu \vec{r}_n + p_\nu \cos \vec{k}_\nu \vec{r}_n), \quad (7.1)$$

where $\bar{\mu} = \mu/\sqrt{\varepsilon_0 V}$, V is the volume of quantization, p_ν, q_ν are the canonical variables of the harmonic ν , and $\vec{r}_n$ the position vector of the atom n . In this case, we consider the operator V_j from the equation (3.1), of the form [29]:

$$V_j = \sum_\nu (a_{j\nu} p_\nu + b_{j\nu} q_\nu) + \sum_n (A_{jn} \sigma_x^n + B_{jn} \sigma_y^n + C_{jn} \sigma_z^n). \quad (7.2)$$

With the expressions (7.1) and (7.2) we obtain the master equation:

$$\begin{aligned} \frac{d\rho}{dt} = & -\frac{i}{\hbar} [H_0, \rho] - i\frac{\bar{\mu}}{\hbar} \sum_{n\nu} [(\nu\omega q_\nu \sin \vec{k}_\nu \vec{r}_n + p_\nu \cos \vec{k}_\nu \vec{r}_n) \sigma_x^n, \rho] \\ & + \sum_\nu \left\{ i\frac{\lambda_\nu}{2\hbar} ([p_\nu, \rho q_\nu + q_\nu \rho] - [q_\nu, \rho p_\nu + p_\nu \rho]) - \frac{D_{qq\nu}}{\hbar^2} [p_\nu, [p_\nu, \rho]] - \frac{D_{pp\nu}}{\hbar^2} [q_\nu, [q_\nu, \rho]] \right. \\ & + \frac{D_{\rho q\nu}}{\hbar^2} ([q_\nu, [p_\nu, \rho]] + [p_\nu, [q_\nu, \rho]]) \left. \right\} + \sum_n \left\{ -\Lambda_{xy}^n [\sigma_z^n, [\sigma_z^n, \rho]] - \Lambda_{yz}^n [\sigma_x^n, [\sigma_x^n, \rho]] \right. \\ & - \Lambda_{zx}^n [\sigma_y^n, [\sigma_y^n, \rho]] + \Gamma_{xy}^n ([\sigma_x^n, [\sigma_y^n, \rho]] + [\sigma_y^n, [\sigma_x^n, \rho]]) + \Gamma_{yz}^n ([\sigma_y^n, [\sigma_z^n, \rho]] + [\sigma_z^n, [\sigma_y^n, \rho]]) \\ & + \Gamma_{zx}^n ([\sigma_z^n, [\sigma_x^n, \rho]] + [\sigma_x^n, [\sigma_z^n, \rho]]) - \frac{i}{2} [D_x^n ([\sigma_y^n, \sigma_z^n \rho + \rho \sigma_z^n] - [\sigma_z^n, \sigma_y^n \rho + \rho \sigma_y^n]) \\ & + D_y^n ([\sigma_z^n, \sigma_x^n \rho + \rho \sigma_x^n] - [\sigma_x^n, \sigma_z^n \rho + \rho \sigma_z^n]) + D_z^n ([\sigma_x^n, \sigma_y^n \rho + \rho \sigma_y^n] - [\sigma_y^n, \sigma_x^n \rho + \rho \sigma_x^n])] \left. \right\}, \quad (7.3) \end{aligned}$$

where H_0 is the Hamiltonian of the system without interaction, and the phenomenological parameters are defined by the expression:

$$D_{qq\nu} = \frac{\hbar}{2} \sum_j a_{j\nu}^* a_{j\nu}, \quad D_{pp\nu} = \frac{\hbar}{2} \sum_j b_{j\nu}^* b_{j\nu},$$

$$\begin{aligned}
D_{pq\nu} &= -\frac{\hbar}{4} \sum_j (a_{j\nu}^* b_{j\nu} + a_{j\nu} b_{j\nu}^*), \quad \lambda_\nu = \frac{i}{2} \sum_j (a_{j\nu}^* b_{j\nu} - a_{j\nu} b_{j\nu}^*), \\
\Lambda_{xy}^n &= \frac{1}{2\hbar} \sum_j C_{jn}^* C_{jn}, \quad \Lambda_{yz}^n = \frac{1}{2\hbar} \sum_j A_{jn}^* A_{jn}, \quad \Lambda_{zx}^n = \frac{1}{2\hbar} \sum_j B_{jn}^* B_{jn}, \\
\Gamma_{xy}^n &= -\frac{1}{4\hbar} \sum_j (A_{jn}^* B_{jn} + A_{jn} B_{jn}^*), \quad D_z^n = -\frac{i}{2\hbar} \sum_j (A_{jn}^* B_{jn} - A_{jn} B_{jn}^*), \\
\Gamma_{yz}^n &= -\frac{1}{4\hbar} \sum_j (B_{jn}^* C_{jn} + B_{jn} C_{jn}^*), \quad D_x^n = -\frac{i}{2\hbar} \sum_j (B_{jn}^* C_{jn} - B_{jn} C_{jn}^*), \\
\Gamma_{zx}^n &= -\frac{1}{4\hbar} \sum_j (C_{jn}^* A_{jn} + C_{jn} A_{jn}^*), \quad D_y^n = -\frac{i}{2\hbar} \sum_j (C_{jn}^* A_{jn} - C_{jn} A_{jn}^*). \tag{7.4}
\end{aligned}$$

Separating the field from the atomic observables by the "mean-field approximation", for the atomic expectation values we obtain the equations

$$\begin{aligned}
\frac{d}{dt} \langle \sigma_x \rangle + \gamma_\perp' \langle \sigma_x \rangle - (\omega_0 - s) \langle \sigma_y \rangle + \gamma_1 \langle \sigma_z \rangle - D_1 &= 0, \\
\frac{d}{dt} \langle \sigma_y \rangle + (\omega_0 - s) \langle \sigma_x \rangle + \gamma_\perp'' \langle \sigma_y \rangle + (\gamma_2 - \bar{\chi}) \langle \sigma_z \rangle - D_2 &= 0, \\
\frac{d}{dt} \langle \sigma_z \rangle + \gamma_1 \langle \sigma_x \rangle + (\gamma_2 + \bar{\chi}) \langle \sigma_y \rangle + \gamma_\parallel \langle \sigma_z \rangle - D_3 &= 0, \tag{7.5}
\end{aligned}$$

where

$$\begin{aligned}
\gamma_\perp' &= 4(\Lambda_{xy} + \Lambda_{zx}), \quad \gamma_\perp'' = 4(\Lambda_{xy} + \Lambda_{yz}), \quad \gamma_\parallel = 4(\Lambda_{zx} + \Lambda_{yz}), \\
s &= 4\Gamma_{xy}, \quad \gamma_1 = 4\Gamma_{zx}, \quad \gamma_2 = 4\Gamma_{yz}, \quad D_1 = 4D_x, \quad D_2 = 4D_y, \quad D_3 = 4D_z \tag{7.6}
\end{aligned}$$

and

$$\bar{\chi} = \frac{2\bar{\mu}}{\hbar} \sum_\nu \text{Tr}[(\nu\omega q_\nu + p_\nu)\rho] \tag{7.7}$$

is a normalized field variable. From these equations one can obtain the classical Bloch-Feynman equations when $s = \gamma_1 = \gamma_2 = 0$, $D_1 = D_2 = 0$, $\gamma_\perp' = \gamma_\perp''$ [29].

Having in view the equation (3.3), in the master equation (7.3), we distinguish two additional parts describing the interaction with the environment: one part for the harmonic oscillators corresponding to the electromagnetic field and another part for the ensemble of atoms. This latter part describes three processes of the atom interaction with the environment: 1) the friction, described by the parameters Λ or $\gamma_\perp', \gamma_\perp'', \gamma_\parallel$; 2) the diffusion, leading

to zero-point values of the observables, described by the parameters D_i ; 3) the coupling between the observables described by the parameters Γ or s , γ_1, γ_2 . These parameters satisfy fundamental constraints, similar to the expressions (3.4) and (3.24):

$$\begin{aligned}
D_{qq\nu} &\geq 0, \quad D_{pp\nu} \geq 0, \quad D_{pp\nu}D_{qq\nu} - D_{pq\nu}^2 \geq \frac{\hbar^2\lambda_\nu^2}{4}, \\
D_{qq\nu}\Delta_{pp\nu}(t) + D_{pp\nu}\Delta_{qq\nu}(t) - 2D_{pq\nu}\Delta_{pq\nu}(t) &\geq \frac{\hbar^2\lambda_\nu^2}{4}, \\
\Lambda_{xy}^n &\geq 0, \quad \Lambda_{yz}^n \geq 0, \quad \Lambda_{zx}^n \geq 0, \\
\Lambda_{xy}^n\Lambda_{zx}^n - \Gamma_{yz}^{n^2} &\geq \frac{1}{4}D_x^{n^2}, \quad \Lambda_{yz}^n\Lambda_{xy}^n - \Gamma_{zx}^{n^2} \geq \frac{1}{4}D_y^{n^2}, \quad \Lambda_{zx}^n\Lambda_{yz}^n - \Gamma_{xy}^{n^2} \geq \frac{1}{4}D_z^{n^2}, \\
\Lambda_{xy}^n\Delta_{zz}^n(t) + \Lambda_{zx}^n\Delta_{yy}^n(t) - 2\Gamma_{yz}^n\Delta_{yz}^n(t) &\geq D_x^n < \sigma_x^n(t) >, \\
\Lambda_{yz}^n\Delta_{xx}^n(t) + \Lambda_{xy}^n\Delta_{zz}^n(t) - 2\Gamma_{zx}^n\Delta_{zx}^n(t) &\geq D_y^n < \sigma_y^n(t) >, \\
\Lambda_{zx}^n\Delta_{yy}^n(t) + \Lambda_{yz}^n\Delta_{xx}^n(t) - 2\Gamma_{xy}^n\Delta_{xy}^n(t) &\geq D_z^n < \sigma_z^n(t) >, \tag{7.8}
\end{aligned}$$

where $\Delta_{pp}, \Delta_{qq}, \Delta_{xx}, \Delta_{yy}, \Delta_{zz}$ are the variances of the observables and the quantities $\Delta_{pq}, \Delta_{xy}, \Delta_{yz}, \Delta_{zx}$ are expressions of the form:

$$\Delta_{pq} = \langle \frac{pq + qp}{2} \rangle - \langle p \rangle \langle q \rangle. \tag{7.9}$$

Following the model of the geometrical representation originally adapted by Feynman, Vernon and Hellwarth [95], we consider the atomic observables with the "amplitudes" u, v, w , as components of the Bloch vector in a "rotating frame" of the frequency ω :

$$\langle \sigma_x \rangle = u \cos \omega t - v \sin \omega t, \quad \langle \sigma_y \rangle = -u \sin \omega t - v \cos \omega t, \quad \langle \sigma_z \rangle = -w. \tag{7.10}$$

For a system of atoms with the density $\mathbf{N}$, using (7.10) one obtains the following expressions of the macroscopic polarization $S = \mathbf{N} \langle \sigma_x \rangle$ and of the population $N = \mathbf{N} \langle \sigma_z \rangle$:

$$S = \frac{1}{2}(\mathbf{S}e^{-i\omega t} + \mathbf{S}^*e^{i\omega t}), \quad N = -\mathbf{N}w, \tag{7.11}$$

where

$$\mathbf{S} = \mathbf{N}(u - iv). \tag{7.12}$$

With the expressions (7.10)–(7.12), the equations (7.5) become:

$$\frac{d\mathbf{S}}{dt} + (\gamma_\perp + \Delta)\mathbf{S} + \gamma_a\mathbf{S}^*e^{2i\omega t} = (i\chi - \gamma e^{i\omega t} + i\chi^*e^{2i\omega t})N + De^{i\omega t},$$

$$\frac{dN}{dt} + \gamma_{\parallel}(N - N_3) = \frac{1}{2} [(i\chi^* - \gamma^* e^{i\omega t} + i\chi e^{-2i\omega t})\mathbf{S} - (i\chi - \gamma e^{i\omega t} + i\chi^* e^{2i\omega t})\mathbf{S}^*], \quad (7.13)$$

where

$$\begin{aligned} \gamma_{\perp} &= (\gamma'_{\perp} + \gamma''_{\perp})/2, \quad \gamma_a = (\gamma'_{\perp} - \gamma''_{\perp})/2, \quad \Delta = \omega_0 - s - \omega, \\ \gamma &= \gamma_1 + i\gamma_2, \quad D = \mathbf{N}(D_1 + iD_2), \quad N_3 = \mathbf{N}D_3/\gamma_{\parallel}. \end{aligned} \quad (7.14)$$

By neglecting the rapidly varying terms, one obtains the classical optical Bloch equations in the "rotating wave approximation" :

$$\begin{aligned} \frac{d\mathbf{S}}{dt} + (\gamma_{\perp} + i\Delta)\mathbf{S} &= i\chi N, \\ \frac{dN}{dt} + \gamma_{\parallel}(N - N_3) &= \frac{i}{2}(\mathbf{S}\chi^* - \mathbf{S}^*\chi). \end{aligned} \quad (7.15)$$

In order to include the effects of these terms we consider solutions of the form:

$$\begin{aligned} E &= \mathbf{E}_0 + \frac{1}{2}(\mathbf{E}_1 e^{-i\omega t} + \mathbf{E}_1^* e^{i\omega t}) + \frac{1}{2}(\mathbf{E}_2 e^{-2i\omega t} + \mathbf{E}_2^* e^{2i\omega t}) + \dots, \\ S &= \mathbf{S}_0 + \frac{1}{2}(\mathbf{S}_1 e^{-i\omega t} + \mathbf{S}_1^* e^{i\omega t}) + \frac{1}{2}(\mathbf{S}_2 e^{-2i\omega t} + \mathbf{S}_2^* e^{2i\omega t}) + \dots, \\ N &= N_0 + \frac{1}{2}(\mathbf{N}_1 e^{-i\omega t} + \mathbf{N}_1^* e^{i\omega t}) + \frac{1}{2}(\mathbf{N}_2 e^{-2i\omega t} + \mathbf{N}_2^* e^{2i\omega t}) + \dots, \end{aligned} \quad (7.16)$$

where $\mathbf{E}_0, \mathbf{E}_1, \mathbf{E}_2, \dots, \mathbf{S}_0, \mathbf{S}_1, \mathbf{S}_2, \dots, \mathbf{N}_0, \mathbf{N}_1, \mathbf{N}_2, \dots$ are slowly varying functions which are called "amplitudes". For the amplitudes of zeroth and first order we get the following equations:

$$\begin{aligned} \frac{d\mathbf{S}_1}{dt} + (\gamma'_{\perp} + i\Delta)\mathbf{S}_1 &= -\gamma_1 \mathbf{N}_1, \\ \gamma_a \mathbf{S}_1 + \frac{\gamma_1 - i(\gamma_2 - 2\chi_0)}{2} \mathbf{N} + i\chi_1 \mathbf{N}_0 &= 0, \\ \frac{d\mathbf{N}_0}{dt} + \gamma_{\parallel}(\mathbf{N}_0 - N_3) + \gamma_1 \mathbf{S}_0 &= \frac{1}{2}(\mathbf{S}_1 \chi_1^* - \mathbf{S}_1^* \chi_1), \\ \frac{d\mathbf{S}_0}{dt} + \gamma'_{\perp} \mathbf{S}_0 + \gamma_1(N_0 - N_1) &= 0, \end{aligned} \quad (7.17)$$

where $\chi_0 = \mu \mathbf{E}_0 / \hbar$, $\chi_1 = \mu \mathbf{E}_1 / \hbar$, $N_1 = \mathbf{N}D_1/\gamma_1$.

In these equations the new parameter $\gamma_1 \neq 0$ leads to a coupling of the polarization variable $\mathbf{S}_1$ with the electric field variable χ_1 , i.e. to an interaction of the atom with the electric field. In this case one obtains the polarization equation

$$\frac{d\mathbf{S}_1}{dt} + \left(\frac{\gamma''_{\perp} - i\gamma'_{\perp}(\gamma_2 - 2\chi_0)/\gamma_1}{1 - i(\gamma_2 - 2\chi_0)/\gamma_1} + i\Delta \right) \mathbf{S}_1 = \frac{2i\chi_1 \mathbf{N}_0}{1 - i(\gamma_2 - 2\chi_0)/\gamma_1}, \quad (7.18)$$

which can be compared with the conventional Bloch polarization equation (7.15). We see that this equation describes new physical phenomena. For instance, in the steady state, with the notations

$$\begin{aligned}\Gamma &= \frac{\gamma_2 - 2\chi_0}{\gamma_1}, \quad \zeta = \frac{\gamma'_\perp}{\gamma''_\perp} \Gamma, \quad \gamma'_\parallel = \gamma_\parallel \left(1 - \frac{\gamma_1^2}{\gamma'_\perp \gamma_\parallel}\right), \quad \chi_s^2 = \frac{\gamma'_\parallel \gamma''_\perp}{2}, \\ N^e &= \frac{N_3 - \frac{\gamma_1 N_1}{\gamma'_\perp}}{1 - \frac{\gamma_1^2}{\gamma'_\perp \gamma_\parallel}}, \quad \delta = \frac{\Delta}{\gamma''_\perp}, \quad \varepsilon = \frac{\chi_1}{\chi_s},\end{aligned}\tag{7.19}$$

one obtains the polarization amplitude

$$\mathbf{S}_1 = 2N^e \frac{\chi_s}{\gamma''_\perp} \cdot \frac{[\delta - \zeta + i(1 + \Gamma\delta)]\varepsilon}{(1 + \Gamma\delta)^2 + (\delta - \zeta)^2 + (1 + \Gamma\delta)|\varepsilon|^2}.\tag{7.20}$$

For the slowly varying amplitudes χ_{1+} and χ_{1-} of the field variables,

$$\chi_1 = \chi_{1+} e^{ikz} + \chi_{1-} e^{-ikz},\tag{7.21}$$

the Maxwell equations take the form [96]:

$$\begin{aligned}\frac{d\chi_{1+}}{dz} &= i \frac{\mu g}{c\hbar} \mathbf{S}_{1+}, \\ \frac{d\chi_{1-}}{dz} &= -i \frac{\mu g}{c\hbar} \mathbf{S}_{1-}, \\ \chi_0 &= -\frac{\mu^2}{\varepsilon_0 \hbar} \mathbf{S}_0,\end{aligned}\tag{7.22}$$

where $g = \omega\mu/2\varepsilon_0$ is the coupling coefficients and $\mathbf{S}_{1+}, \mathbf{S}_{1-}$ are the Fourier transforms of the polarization $\mathbf{S}_1$ as a function of the coordinate z :

$$\mathbf{S}_{1+} = \frac{k}{2\pi} \int_{-\pi}^{\pi} \mathbf{S}_1(z) e^{-ikz} dz, \quad \mathbf{S}_{1-} = \frac{k}{2\pi} \int_{-\pi}^{\pi} \mathbf{S}_1(z) e^{ikz} dz.\tag{7.23}$$

Eqs. (7.22) with the expression (7.20) describe the propagation of an electromagnetic plane wave, taking into account the self-reflection effect [97]. Neglecting the counter-propagating wave due to the self-reflection, one obtains the absorption coefficient

$$\alpha = -\frac{1}{|\varepsilon|} \frac{d|\varepsilon|}{dz} = \frac{\alpha_0}{2} \frac{1 + \Gamma\delta}{(1 + \Gamma\delta)^2 + (\delta - \zeta)^2 + (1 + \Gamma\delta)|\varepsilon|^2}\tag{7.24}$$

and the corresponding dephasing

$$\frac{d\theta}{dz} = -\frac{\delta - \zeta}{1 + \Gamma\delta}\alpha, \quad (7.25)$$

where $\alpha_0 = 4\mu g N^e / e\hbar\gamma_{\perp}^{\parallel}$.

According to Eq. (7.24) the absorption coefficient α becomes negative when the electric field intensity $|\varepsilon|^2$ is sufficiently small for values of the atomic detuning δ with $1 + \Gamma\delta < 0$. In that case the electromagnetic wave is amplified, taking energy from the environment. At the same time, the difference between the population of the ground state and the population of the excited state,

$$N_0 = \frac{N^e}{1 + \frac{1+\Gamma\delta}{(\delta-\zeta)^2 + (1+\Gamma\delta)^2} \frac{|\chi_1|^2}{\chi_s^2}} \quad (7.26)$$

becomes larger than its equilibrium value N^e . When the ensemble of atoms has a smaller temperature than the environment, the energy passes from the environment to the atomic system and finally to the electromagnetic field. The absorption coefficient asymmetry with the atomic detuning δ , can be observed in the experimental data of Sandle and Gallagher [27,98]. The laser radiation absorption by Na atoms in a buffer gas of Ar atoms has been studied by McCartan and Farr [99]. They varied the pressure of the buffer gas and measured the spectral line width w and the resonance frequency shift $\Delta\omega_0$. Using the photoluminescence method [100], we obtain the following expression of the spectral line width

$$W = 2\pi\Delta\omega^2 \frac{\alpha}{\int \alpha d\omega}, \quad (7.27)$$

while the resonance frequency shift becomes

$$\Delta\omega_0 = \gamma_{\perp}^{\prime\prime}\xi. \quad (7.28)$$

Integrating the expression (7.24) one obtains

$$W = 2\gamma_{\perp}^{\prime\prime}(1 + \Gamma\xi). \quad (7.29)$$

By taking into account (7.28), this expression becomes

$$W = 2\Delta\omega_0 \frac{1 + \Gamma\xi}{\xi} \simeq 2\Delta\omega_0. \quad (7.30)$$

Consequently the spectral line width W is proportional to the line frequency shift $\Delta\omega_0$ which is in agreement with the experimental results of McCartan and Farr [99].

8. Summary

The Lindblad theory provides a selfconsistent treatment of damping as a possible extension of quantum mechanics to open systems. In the present review first we studied the damped quantum oscillator by using the Schrödinger and Heisenberg representations. According to this theory we have calculated the damping of the expectation values of coordinate and momentum and the variances as functions of time. The resulting time dependence of the expectation values yields an exponential damping. Then we have shown that the quasiprobability distributions can be used to solve the problem of dissipation for the harmonic oscillator. From the master equation of the damped quantum oscillator we have derived the corresponding Fokker-Planck equations in the Glauber P , the antinormal ordering Q and the Wigner W representations and have made a comparative study of these quasiprobability distributions. The Fokker-Planck equations we obtained describe an Ornstein-Uhlenbeck process. We have proven that the variances found from the Fokker-Planck equations in these representations are the same. We have solved these equations in the steady state and showed that the Glauber P function (when it exists), the Q and the Wigner W functions are two-dimensional Gaussians with different widths. We have also calculated the time evolution of the density matrix. For this purpose we applied the method of the generating function of the density matrix. In this case the density matrix can be obtained by taking partial derivatives of the generating function. The generating function depends on a set of time-dependent coefficients which are calculated as solutions of linear differential equations of first order. Depending on the initial conditions for these coefficients, the density matrix evolves differently in time. When the asymptotic state is a Gibbs state in the case of a thermal bath, a Bose-Einstein distribution results as density matrix. Also for the case that the initial density matrix is chosen as a Glauber packet, a simple analytical expression for the density matrix has been derived. The density matrix can be used in various physical applications where a bosonic degree of freedom moving in a harmonic oscillator potential is damped. For example, one needs to determine nondiagonal transition elements of the density matrix, for an oscillator perturbed by a weak electromagnetic field in addition to its coupling to a heat bath. The density matrix

can also be derived from the solution of the Fokker-Planck equation for the coherent state representation.

The theory was applied to the damping of the charge equilibration in deep inelastic collisions of heavy ions. The comparison of theoretical results with experimental data shows that the overdamped solution succeeded to describe these experimental data.

We have also shown the Hamiltonian of the proton and neutron asymmetry degrees of freedom in deep inelastic collisions as an example for two coupled and damped oscillators. The usual limitation of the Lindblad theory is that the damping time is long compared with the characteristic time of the oscillators. This condition is not too well satisfied in deep inelastic collisions of nuclei, where the time scale is of the order of the relaxation time. Therefore, in these applications we consider the Lindblad theory as an axiomatic procedure for describing the dissipation processes and accept its parameters as free quantities, fitted to the experimental data.

We obtained simple models for the dissipation of the angular momentum, by using the generators of the Lorentz group as openness operators of the system. For given values of the diffusion and friction coefficients we obtained an exponential damping of the angular momentum, in agreement with the situation encountered in heavy ion collisions. If the opening of the system fulfils rotational symmetry and the Hamiltonian is a positive function of L^2 and N^2 , (square of generators of Lorentz group), then the Lindblad master equation describes a physical system coupled to a reservoir of infinite temperature, which increases the energy and angular momentum exponentially.

We calculated the tunneling spectrum as a function of some barrier characteristics: the tunneling operator, the overlap integral, the transition elements of the coordinate and momentum. Besides Gamow's tunneling process with energy conservation, additional terms with energy transfer to or from environment are obtained. For low values of temperature, the whole spectrum is situated at energies smaller than the Q -value; for higher enough values of temperature, transitions at energies higher than the Q -value appear. Generally, we shown that dissipation stimulates the tunneling process and leads to an exotic decay law.

Using Lindblad's theory of open systems, for the resonant atom-field interaction, we found new optical equations. We showed that the interaction with the environment

consists not only in a decay, but also in a coupling of the atomic observables. This coupling has experimental evidence in the absorption spectrum of the laser radiation and in optical bistability.

At the same time, this phenomenon leads to a remarkable physical effect of energy transfer from the dissipative environment, which is cooling, to the coherent electromagnetic field propagating through it.

Recently we assist to a revival of interest in quantum brownian motion as a paradigm of quantum open systems. There are many motivations. The possibility of preparing systems in macroscopic quantum states led to the problems of dissipation in tunneling and of loss of quantum coherence (decoherence). These problems are intimately related to the issue of quantum-to-classical transition. All of them point the necessity of a better understanding of open quantum systems and all requires the extension of the model of quantum brownian motion. Our results allow such extensions and also explain some earlier observations. For a first comment of the result (3.4),*iii*) see Sec. 4.2 of Ref. [101]. The fact that the Schrödinger equation corresponding to the Lindblad equation is nonlinear was obtained in Ref. [17] and was quoted in Ref. [102]. The approach used in Ref. [17] to generate the Dekker master equation was applied in Ref. [93] to generate a family of master equations for local quantum dissipation. New developments of this kind were also obtained in Ref. [103].

References

1. R. W. Hasse, J. Math. Phys. **16** (1975) 2005
2. E. B. Davies, Quantum Theory of Open Systems, Academic Press, New York, 1976
3. J. Messer, Acta Phys. Austriaca **58** (1979) 75
4. H. Spohn, Rev. Mod. Phys. **52** (1980) 569
5. H. Dekker, Phys. Rep. **80** (1981) 1
6. P. Exner, Open Quantum Systems and Feynman Integrals, Reidel, Dordrecht, 1985
7. K. H. Li, Phys. Rep. **134** (1986) 1
8. R. Haake, Springer Tracts in Mod. Phys. **66** (1973) 98
9. R. S. Ingarden and A. Kossakowski, Ann. Phys. (N.Y.) **89** (1975) 451
10. V. Gorini and A. Kossakowski, J. Math. Phys. **17** (1976) 1298
11. A. Kossakowski, Rep. Math. Phys. **3** (1972) 247

12. A. Kossakowski, Bull. Acad. Polon. Sci. Math. Astron. Phys. **20** (1972) 1021
13. R. S. Ingarden, Acta Phys. Polonica A **43** (1973) 1
14. G. Lindblad, Commun. Math. Phys. **48** (1976) 119
15. V. Gorini, A. Frigerio, M. Verri, A. Kossakowski and E. C. G. Sudarshan, Rep. Math. Phys. **13** (1978) 149
16. G. Lindblad, Rep. Math. Phys. **10** (1976) 393
17. A. Sandulescu and H. Scutaru, Ann. Phys. (N.Y.) **173** (1987) 277
18. A. Isar, A. Sandulescu and W. Scheid, J. Phys. G - Nucl. Part. Phys. **17** (1991) 385
19. A. Isar, W. Scheid and A. Sandulescu, J. Math. Phys. **32** (1991) 2128
20. A. Isar, A. Sandulescu and W. Scheid, J. Math. Phys. **34** (1993) 3887
21. S. Jang, Nucl. Phys. A **499** (1989) 250
22. A. Pop, A. Sandulescu, H. Scutaru and W. Greiner, Z. Phys. A **329** (1988) 357
23. A. Sandulescu, H. Scutaru and W. Scheid, J. Phys. A - Math. Gen. **20** (1987) 2121
24. A. Isar, A. Sandulescu, H. Scutaru and W. Scheid, Nuovo Cimento A **103** (1990) 413
25. A. Isar, A. Sandulescu and W. Scheid, Int. J. Mod. Phys. A **5** (1990) 1773
26. E. Stefanescu, A. Sandulescu and W. Greiner, Int. J. Mod. Phys. E **2** (1993) 233
27. M. Razavi and A. Pimpale, Phys. Rep. **168** (1988) 305
28. H. Dekker, Phys. Rev. A **38** (1988) 6351
29. A. Sandulescu and E. Stefanescu, Physica A **161** (1989) 525
30. G. C. Emch, Algebraic Methods in Statistical Mechanics and Quantum Field Theory, Wiley, New York, 1972
31. V. Gorini, A. Kossakowski and E. C. G. Sudarshan, J. Math. Phys. **17** (1976) 821
32. P. Talkner, Ann. Phys. (N.Y.) **167** (1986) 390
33. H. Dekker, Phys. Rev. A **16** (1977) 2126
34. H. Dekker, Phys. Lett. A **74** (1979) 15
35. H. Dekker, Phys. Lett. A **80** (1980) 369
36. H. Dekker and M. C. Valsakumar, Phys. Lett. A **104** (1984) 67
37. E. Merzbacher, Quantum Mechanics, Wiley, New York, 1970
38. H. Hofmann and P. J. Siemens, Nucl. Phys. A **275** (1977) 464
39. H. Hofmann, C. Grégoire, R. Lucas and C. Ngô, Z. Phys. A **293** (1979) 229
40. R. W. Hasse, Nucl. Phys. A **318** (1979) 480

41. E. M. Spina and H. A. Weidenmüller, Nucl. Phys. A **425** (1984) 354
42. C. W. Gardiner and M. J. Collet, Phys. Rev. A **31** (1985) 3761
43. T. A. B. Kennedy and D. F. Walls, Phys. Rev. A **37** (1988) 152
44. G. S. Agarwal, Phys. Rev. **178** (1969) 2025
45. G. S. Agarwal, Phys. Rev. A **4** (1971) 739
46. S. Dattagupta, Phys. Rev. A **30** (1984) 1525
47. C. M. Savage and D. F. Walls, Phys. Rev. A **32** (1985) 2316
48. N. Lu, S. Y. Zhu and G. S. Agarwal, Phys. Rev. A **40** (1989) 258
49. S. Jang and C. Yannouleas, Nucl. Phys. A **460** (1986) 201
50. S.Chaturvedi, P.D.Drummond and D.F.Walls, J. Phys. A - Math. Gen. **10** (1977) L187
51. P. D. Drummond and C. W. Gardiner, J. Phys. A - Math. Gen. **13** (1980) 2353
52. P. D. Drummond, C. W. Gardiner and D. F. Walls, Phys. Rev. A **24** (1981) 914
53. M. Hillery, R. F. O 'Connell, M. O. Scully and E. P. Wigner, Phys. Rep. **106** (1984) 121
54. E. P. Wigner, Phys. Rev. **40** (1932) 749
55. E. J. Glauber, Phys. Rev. **131** (1963) 2766
56. E. J. Glauber, Phys. Rev. Lett. **10** (1963) 84
57. E. C. G. Sudarshan, Phys. Rev. Lett. **10** (1963) 277
58. W. Weidlich, H. Risken and H. Haken, Z. Phys. **204** (1967) 223
59. M. Lax and W. H. Louisell, IEEE J. Q. Electron. **3** (1967) 47
60. K. E. Cahill and R. J. Glauber, Phys. Rev. A **177** (1969) 1882
61. R. J. Glauber, in Laser Handbook, ed. by F. T. Arecchi and E. O. Schultz-Dubois, North-Holland, Amsterdam, 1972
62. J. R. Klauder and E. C. G. Sudarshan, Fundamentals of Quantum Optics, Benjamin, New York, 1968
63. K. E. Cahill, Phys. Rev. **180** (1969) 1239
64. K. E. Cahill, Phys. Rev. **180** (1969) 1244
65. W. H. Louisell, Quantum Statistical Properties of Radiation, Wiley, New York, 1973
66. H. Haken, Rev. Mod. Phys. **47** (1975) 67
67. C. W. Gardiner, Handbook of Stochastic Methods, Springer, Berlin, 1982

68. G. E. Uhlenbeck and L. S. Ornstein, Phys. Rev. **36** (1930) 823
69. M. C. Wang and G. E. Uhlenbeck, Rev. Mod. Phys. **17** (1945) 323
70. H. Weyl, The Theory of Groups and Quantum Mechanics, Dover, New York, 1931
71. H. Freiesleben and J. B. Kratz, Phys. Rep. **106** (1984) 1
72. W.U.Schröder and J.R.Huizenga, Treatise on Heavy-Ion Science, vol.2, Plenum, New York, 1984
73. J.A.Maruhn, W.Greiner and W.Scheid, Heavy Ion Collisions, vol.2, North-Holland, Amsterdam, 1980
74. A. Sandulescu, M. Petrovici, A. Pop, M. S. Popa, T. Hahn, K. H. Ziegenhain and W. Greiner, J. Phys. G - Nucl. Part. Phys. **7** (1981) L55
75. E. S. Hernandez, W. D. Myers, J. Randrup and B. Remaud, Nucl. Phys. A **361** (1981) 483
76. H. Breuer, A. C. Mignerey, V. E. Viola, K. L. Wolf, J. R. Birkelund, D. Hilscher, J. R. Huizenga, W. V. Schröder and W. W. Wilde, Phys. Rev. C **28** (1983) 1080
77. G. Wolschin, Nukleonika **22** (1977) 1165
78. G. Wolschin, Fizika **9** Suppl. 4 (1977) 415
79. C. Riedel, G. Wolschin and W. Nörenberg, Z. Phys. A **290** (1979) 47
80. R. K. Gupta, M. Münchow, A. Sandulescu and W. Scheid, J. Phys. G. **10** (1984) 209
81. D. H. E. Gross and K. H. Hartmann, Phys. Rev. C **24** (1981) 2526
82. W. V. Schröder, J. R. Huizenga and J. Randrup, Phys. Lett. B **98** (1981) 355
83. J. R. Birkelund, H. Freiesleben, J. R. Huizehga, W. V. Schröder, W. W. Wilcke, K. L. Wolf, J. P. Unik and V. E. Viola, Phys. Rev. C **26** (1982) 1984
84. A. C. Merchant and W. Nörenberg, Z. Phys. A **308** (1982) 315
85. V. V. Dodonov and O. V. Manko, Physica A **130** (1985) 353
86. A. A. Belavin, B. Ya. Zel'dovich, A. M. Perelomov and V. S. Popov, Z. Eksp. Teor. Fiz. **56** (1969) 264
87. Wu-Ki Tung, Group Theory in Physics, World Scientific, Singapore, 1985
88. A. Isar, A. Sandulescu and W. Scheid, Rev. Roum. Phys. **35** (1990) 13
89. G. Gamow, Z. Phys. **51** (1928) 204
90. A. O. Caldeira and A. J. Leggett, Ann. Phys. (N.Y.) **149** (1983) 374
91. K. Fujikawa et al., Phys. Rev. Lett. **68** (1992) 1093

92. E. G. Harris, Phys. Rev. A **48** (1993) 995
93. M. R. Gallis, Phys. Rev. A **48** (1993) 1028
94. D. M. Brink, private comm.
95. R. P. Feynman, F. L. Vernon Jr. and R. W. Hellwarth, J. Appl. Phys. **28** (1957) 49
96. H. J. Carmichael, Optica Acta **27** (1980) 147
97. L. Roso-Franco, J. Opt. Soc. Amer. **4** (1987) 1878
98. W. J. Sandle and A. Gallagher, Phys. Rev. A **24** (1981) 2017
99. D. G. McCartan and J. M. Farr, J. Phys. B **9** (1976) 985
100. R. H. Chatham, A. Gallagher and E. L. Lewis, J. Phys. B **13** (1980) L7
101. V. V. Dodonov, V. I. Man'ko, O. V. Man'ko, Quantum nonstationary oscillator, Trudy FIAN, vol. 191, Nauka, Moskow, 1989
102. A. Iannussis and R. Mignani, Physica A **152** (1988) 469
103. N. V. Antonenko, S. P. Ivanova, R. V. Jolos and W. Scheid, J. Phys. G: Nucl. Part. Phys. **20** (1994) 1447